FOREWORD

INTRODUCTION

BENZOYL PEROXIDE

CAS N[•]: 94-36-0

SIDnitial Assessment Report

For

SIAM 15

(Boston, USA, 22-25 October 2002)

- 1. Chemical Name: Benzoyl Peroxide
- **2. CAS Number:** 94-36-0
- **3. Sponsor Country:** Republic of Korea

National SIDS Contact Point in Sponsor Country : Dr. Hyun-Mi Kim National Institute of Environmental Research, Gyeongseo-dong, Seo-gu, Incheon 404-170, Korea Tel : 82-32-560-7124 Fax : 82-32-568-2037 E-mail : hmikim@me.go.kr

4. Shared Partnership with:

5. Roles/Responsibilities of the Partners:

- Name of industry sponsor /consortium
- Process used
- 6. Sponsorship History
- How was the chemical or category brought into the OECD HPV Chemicals Programme ?
- 7. Review Process Prior to the SIAM:
- 8. Quality check process: No testing ()

This substance is sponsored by Korea. The assessment process was started in 1999.

Testing (x):

Algae Growth Inhibition Test (TG 201), Daphnia Acute Immobilization Test (TG 202), Acute Toxicity of Fish (TG 203), Acute Toxicity Test of Earthworm (TG 207), Acute Oral Toxicity (TG 401), Combined Repeated Dose Toxicity with the Reproduction/Developmental Toxicity Screening Test (TG 422), Bacterial Reverse Mutation Test (TG 471), Micronucleus test (TG 474)

- 9. Date of Submission:
- **10. Date of last Update:**
- 11. Comments:

Revised : 2002

SIDS INITIAL ASSESSMENT PROFILE

CAS No.	94-36-0	
Chemical Name	Benzoyl peroxide	
Structural Formula	C ₁₄ H ₁₀ O ₄	

SUMMARY CONCLUSIONS OF THE SIAR

Human Health

The dry chemical becomes explosive above 105°C and when subjected to impact or friction.

The acute oral toxicity of benzoyl peroxide is very low: LD50 >2,000 mg/kg bw in mice, and 5,000 mg/kg bw in rats. No deaths occurred in male rats following inhalation of 24.3 mg/L. Visible effects included eye squint, dyspnea, salivation, lacrimation, erythema and changes of respiratory rates and motor activity.

Benzoyl peroxide was slightly irritating to skins in 24 hr-patch tests. Benzoyl peroxide was not irritating to the eyes of rabbits if washed out within 5 minutes after instillation, however, if the chemical was not washed out until 24 hours later, it proved to be irritating.

Positive results from sensitisation tests in guinea pigs and mice, and from a maximization test in human volunteers, indicate that benzoyl peroxide is a skin sensitizer.

In the combined repeated dose and reproduction/developmental toxicity study (OECD TG 422), benzoyl peroxide did not produce hematological or biochemical adverse effects. Repeated administration by oral gavage up to 1,000 mg/kg bw/day for 29 days resulted in decreased weights of testes and epididymis in male rats. The NOAEL for repeated dose toxicity was 500 mg/kg bw/day.

This substance did not cause gene mutation in bacteria (OECD TG 471 & 472) and *in vitro* chromosomal aberration in CHL (Chinese Hamster Lung) cells. An *in vivo* mammalian erythrocytes micronucleus test (OECD TG 474) produced negative result. The available evidence supports the conclusion that benzoyl peroxide is not a mutagen.

There is no evidence to suggest that benzoyl peroxide is a carcinogen. However, there is some evidence from non-guidelines studies that benzoyl peroxide is a skin tumour promoter.

In the combined repeated dose and reproduction/developmental toxicity study [OECD TG 422], no treatment-related changes in precoital time, rate of copulation, fertility and gestation were noted in any treated group. Adverse effects were shown at the highest dose of 1,000 mg/kg

bw/day in parental male rats with the reduction of reproductive organ weight and slight testes degeneration. In parental female rats, no adverse effects were observed during the test period. The NOAEL for reproduction toxicity in male rats was 500 mg/kg bw/day. In the offspring, the only effect seen was that body weight gain of pups at dose of 1,000 mg/kg bw/day was significantly decreased. The NOAEL for developmental toxicity was 500 mg/kg bw/day.

Environment

Benzoyl peroxide is commercially produced as a white granule with purities ranging from 22 to 95%. It has a melting point of 104 -106 °C, vapor pressure of 0.00929 Pa, solubility of 9.1 mg/L in water at 25 °C, and log P_{ow} of 3.43 at 25 °C. For indirect photolysis in the atmosphere, the half-life is estimated to be 3 days with the AOPWIN model. The substance is readily biodegradable (OECD TG 301C: 83% by BOD after 21 days) and hydrolyses rapidly in water [OECD TG 111] with a half-life of 11.87 hrs at pH 4.0 and 5.20 hr at pH 7.0 at 25 °C. The main hydrolysis product of benzoyl peroxide is benzoic acid (a SIDS assessment of benzoic acid is available: CAS No. 65-85-0). The estimated BCF of 92 with the BCFWIN model suggests that the chemical has a low potential for bioaccumulation.

The following studies for aquatic organisms are available:

Green algae (*Selenastrum capricornutum*): 72 hr- E_bC_{50} is 0.07 mg/L (biomass) and 0.44 mg/L (growth rate). Invertebrates (*Daphnia magna*): 48 hr- EC_{50} is 0.07 mg/L. Fish (*Oryzias latipes*): 96 hr- LC_{50} is 0.24 mg/L. Microorganism (*activated sludge*): 30 min.- EC_{50} is 35 mg/L.

The toxicity observed is assumed to be due to benzoyl peroxide rather than benzoic acid, which shows much lower toxicity to aquatic organisms. One can assume that effects occur before hydrolysis takes place.

A generic fugacity model (Mackay level III) was used for environmental fate estimation. If the most realistic emission pattern to water is assumed then the substance will remain in the aquatic compartment.

Exposure

In Korea, the total production volume of benzoyl peroxide was 1,357 tonnes in 2001, and the chemical is produced in only one company. The amounts of import and export were estimated as 268 and 293 tonnes/year, respectively. 75% benzoyl peroxide is mainly used in the manufacture of expandable styrene polymer and other resins as initiators of polymerization. Benzoyl peroxide has also been used in the treatment of acne vulgaris and the medical product contains mainly 5-10 % benzoyl peroxide. A very small portion of benzoyl peroxide is used as flour bleaching agent.

The Danish, Norwegian and Swedish product register indicates that this substance is used in adhesives, cosmetics, dental products, process regulators, fillers, construction materials and paints. These products may contain 2-80 % of the substance depending upon the product.

The major routes of occupational exposure are inhalation and dermal. Limited data on exposure are available. At a production facility monitoring its workplace for the worker exposure annually, the concentration of airborne aerosols at personal sampling has been less than 1 mg/m^3 .

RECOMMENDATION

The chemical is a candidate for further work.

RATIONALE FOR THE RECOMMENDATION AND NATURE OF FURTHER WORK RECOMMENDED

<u>Human Health</u>: The chemical possesses properties indicating hazards to human health (sensitisation, effect on testes weight, fetal body weight and skin tumour promotion activity) and is a candidate for further work, i.e. exposure assessment, and if considered necessary, risk assessment.

<u>Environment</u>: It is expected that the possibility of any environmental releases of benzoyl peroxide is low in the sponsor country. However, this substance is a candidate for further work, even if it hydrolyses rapidly and has a low bioaccumulation potential. The substance shows high acute toxicity to aquatic organisms and some information indicates wide-dispersive use of this substance. This could lead to local concern for the aquatic environment and therefore environmental exposure assessment is recommended.

SIDS Initial Assessment Report

1 IDENTITY

1.1 Identification of the Substance

CAS Number:	94-36-0
IUPAC Name:	Benzoyl peroxide
Molecular Formula:	$C_{14}H_{10}O_{4}$
Structural Formula:	$(C_6H_5CO)_2O_2$



Molecular Weight: 242.24 Synonyms: Benzoic acid, peroxide Dibenzoyl peroxide Benzoyl superoxide Benzoperoxide Benzovl peroxide, remainder

Benzoyl peroxide, remainder water Lucidol

Benzoyl peroxide (CAS No. 94-36-0) is commercially produced as a white granule with purities ranging from 22 to 99 %. It has peroxy bond in the molecule. Its molecular weight is 242 g/mol. Physical-chemical properties and characteristics are as follow :

1.2 Purity/Impurities/Additives

 $> 97.3 \ \text{W/W}$

1.3 Physico-Chemical properties

Table 1	Summary	of physico-c	hemical	properties
---------	---------	--------------	---------	------------

Property	Value
Melting point	104 - 106 °C
Boiling point	explosive above 105 °C
Relative density	1.33 g/cm ³ at 25 °C
Vapour pressure	0.00929 Pa at 25°C
Water solubility	9.1 mg/L at 25 °C
Partition coefficient n-octanol/water (log value)	3.43 at 25 °C

• Classification in member countries

At present, benzoyl peroxide is not classified as a toxic chemical in the Toxic Chemicals Control Act of the Republic of Korea, but it is in the process of being re-classified as a Toxic Chemical because of its high aquatic toxicity.

2 GENERAL INFORMATION ON EXPOSURE

2.1 **Production Volumes and Use Pattern**

In Korea, the total production volume of benzoyl peroxide was 1,357 tonnes in 2001 and the chemical is produced by only one company. The amounts of import and export were estimated to be 268 and 293 tonnes/year, respectively. Benzoyl peroxide is produced through the reaction of benzoyl chloride with hydrogen peroxide in alkaline solution. Benzoyl peroxide is commercialized as a solid (granules) with varying purities (22 %, 75 % or 95 %). The most often used industrial concentration is 75 %. The substance is mainly used as an initiator for polymerization in the manufacture of expandable styrene polymer and other resins. The world-wide production volume of this chemical is not known. The following table shows the general use pattern of benzoyl peroxide in Korea (Table 1).

Table 1. Use pattern of benzoyl peroxide in Korea

Use	Percentage
Production of expandable styrene (styropor) in the polymer industry	87.9
Production of sizing agent for the textile industry	5.8
Manufacture of acrylic resin for paint products	5.5
Flour bleaching agent in food processing	0.1
Other uses (adhesive, other resins, etc.)	0.2

Benzoyl peroxide has also been used in the treatment of acne vulgaris. All of the medicines containing benzoyl peroxide are imported into Korea with various concentrations as 2.5, 4, 5 and 10 %. Most available acne treatment products contain 5 - 10 % benzoyl peroxide in gel or lotion type.

2.2 Environmental Exposure and Fate

2.2.1 Sources of Environmental Exposure

Benzoyl peroxide hydrolyses to more than 90 % in water after 5 days at 50 °C. Its half-life at 25 °C is 11.9 hr at pH 4 and 5.2 hr at pH 7 (OECD TG 111, NIER, Korea, 2001a). Hydrolysis may be an important environmental fate process of benzoyl peroxide. Benzoyl peroxide is expected to react with hydroxyl radicals in the atmosphere. The estimated half-life is about 3 days (NIER, Korea, 2002a). The substance is readily biodegradable (OECD TG 301C : 83 % by BOD after 21 days). This substance easily passed a ready biodegradation test due to rapid hydrolysis of the parent substance, followed by rapid biodegradation of benzoic acid, which is the hydrolysis product. ABCF of 92, estimated with the BCFWIN model based on a logPow of 3.43, suggests that the chemical has a low potential for bioaccumulation. (Chem. Inspect Test Inst, Japan, 1992 ; Lyman, W.J., et al, 1990 ; OECD, 2000). If it is released into the soil, benzoyl peroxide is expected to have a low mobility in soil based on an estimated Koc value of 1,800 (TOXNET, 2002 ; Lyman, W.J., et al., 1990). Benzoyl peroxide is expected to volatilize from moist soil surfaces based on an estimated Henry's Law constant of 3.5 x 10-6 atm-cu m/mole (Meylan, W.M., 1991). A generic fugacity

model (Mackay level III) is used for the environmental fate estimation. The result is presented in table 2. Benzoyl peroxide is mainly distributed to soil (99.9 %).

Releases to air, water and sediment are negligible. If the substance is emitted to air, it will mainly partition to soil (98.9 %) and if it is released to water, it will mainly remain in water (85.0 %). If it is released to soil, it will mainly remain in soil (99.9 %). (NIER, Korea, 2002a)

Compartment	Release 100 % to air	Release 100 % to water	Release 100 % to soil	All three (%)
Air	1.08	0.068	0.007	0.04
Water	0.01	85.0	0.0068	0.02
Soil	98.9	6.24	99.999	99.9
Sediment	0.001	8.65	0.0007	0.002

Table 2. Environmental Distribution of Benzoyl peroxide using the Fugacity Model (Mackay level III)

Based on the chemical structure, the main hydrolysis product of benzoyl peroxide is benzoic acid. Benzoic acid is mainly distributed to soil (64.2 %) and water (34.8 %) according to a fugacity level III model. No hydrolysis is expected at a pH range of 4 - 11. The substance is readily biodegradable (> 90 % after 28 days) and has a low potential for bioaccumulation (see corresponding initial assessment for benzoic acid, CAS No 65-85-0).

2.3 Human Exposure

2.3.1 Occupational Exposure

Benzoyl peroxide is produced by chemical reaction in a closed reactor. Potential exposure to benzoyl peroxide is limited to the packing process at the production facility. This chemical is used mainly as an initiator for polymerization in which the exposure to this chemical is very limited. The major routes of occupational exposure are inhalation and dermal. The occupational exposure limit of this chemical is set at 5 mg/m³ TWA in Korea. Limited data on exposure are available. Annual monitoring of the concentration of benzoyl peroxide at the workplace of the production facility has shown concentrations in airborne aerosols (personal sampling) of less than 1 mg/m³ (Hansol Chem. Co., 2002).

2.3.2 Consumer Exposure

A very small portion of benzoyl peroxide is used as a flour bleaching agent. The purity of benzoyl peroxide used in the food product is usually less than 22 %. There is no information on actual benzoyl peroxide content in food products. Furthermore, it is well known that benzoyl peroxide is used in the treatment of acne vulgaris and most medical products contain 5 - 10 % benzoyl peroxide.

3 HUMAN HEALTH HAZARDS

3.1 Effects on Human Health

3.1.1. Toxicokinetics, Metabolism and Distribution

Studies in Animals and Humans

In vitro and In vivo Studies

Benzoyl peroxide is rapidly converted to benzoic acid in the skin of both animals and humans. Absorption of the benzoic acid occurs as benzoate through the blood vessels in the dermis. This metabolite enters the blood circulation and is rapidly cleared through the kidney (Nacht, S. *et al*, 1981; Yeung, D., *et al*, 1983; Morsches, B. and Holzmann, H., 1982; Wepierre, J., *et al*, 1986). Topically applied benzoyl peroxide penetrates unchanged through the stratum corneum or follicular openings of excised human skin and is converted metabolically to benzoic acid within the skin (Nacht, S. *et al*, 1981). A study in rhesus monkeys *in vivo* showed that this benzoic acid is systemically absorbed as benzoate and rapidly excreted in the urine in an unchanged form, without being conjugated to hippuric acid, as would be predicted to occur following oral administration.

In vitro and *in vivo* assessments of percutaneous penetration and metabolite analysis of benzoyl peroxide were performed using human skin and in 5 patient with leg ulcers, respectively (Morsches, B., *et al*, 1982). The *in vitro* study showed that benzoyl peroxide absorbed by the skin was converted to benzoic acid in the dermis. The metabolite penetrating through the skin was benzoic acid only. Also in patients treated with benzoyl peroxide, no benzoyl peroxide could be detected in the serum. The authors suggested that benzoyl peroxide was absorbed by the skin, but was systemically absorbed only after transformation to benzoic acid. Therefore, a systemic-toxic effect in local therapy with benzoyl peroxide can be excluded. The rhesus monkey study supports the human study indicating no evidence of parental benzoyl peroxide in the systemic circulation.

3.1.2 Acute Toxicity

Inhalation

In an acute inhalation study, a single group of 10 male rats were placed in a sealed 59.1 liter glass chamber and exposed to a dynamic atmosphere containing the dust of benzoyl peroxide for 4 hrs. All the rats exposed to 24.3 mg/l of 78 % benzoyl peroxide, survived during the 14 days observation period. Observable signs during the 4 hr exposure period were eye squint, dyspnea, salivation, lacrimation, erythema, increased and decreased respiratory rates, and increased and decreased motor activity. A few of the rats showed ocular irritation at 5 days after exposure (Wazeter, F.X., *et al*^(b), 1973). Other acute toxicity studies were reported for rats and mice as shown in table 3.

Route	Species	Туре	Values	References
Oral	Mouse	LD ₅₀	> 2000 mg/kg	NIER, Korea (2001)
Oral	Mouse	LD ₅₀	5700 mg/kg	Sanitariya, G.I. (1936)
Oral	Rat	LD ₅₀	> 5000 mg/kg	Wazeter, F.X. et al (1973)
Oral	Rat	LD ₅₀	> 950 mg/kg	Eastman Kodak Co. (1964)
Oral	Rat	LD ₅₀	7710 mg/kg	Marhold, J.V. (1972)
Inhalation	Rat	LC _{0(4hr)}	24.3 mg/l	Wazeter, F.X. et al (1973)
i.p.	Mouse	LD ₅₀	213 mg/kg	Horgan, V.J. et al (1957)
i.p.	Mouse	LD ₅₀	168 mg/kg	Philpot, J. et al (1959)
i.p.	Mouse	LD_{50}	250 mg/kg	Takahashi, A. (1976)

Table 3. Acute toxicity for rats and mice

Oral

The acute oral toxicity of benzoyl peroxide is likely to be low. In an acute oral toxicity test with ICR mice, a limit test at dose level of 2,000 mg/kg benzoyl peroxide was carried out in a group of 5 males and 5 females using OECD TG 401(NIER^(c), Korea, 2001). There were no deaths in treated animals. No significant findings were observed at necropsy. The only clinical symptom observed in the highest dose group was piloerection. The study concluded that the LD₅₀ value was greater than 2,000 mg/kg for males and females. Similar results were obtained in another acute oral study in rats. No mortality, toxic symptoms or signs were reported at the single dose of 5,000 mg/kg (Wazeter, F.X., *et al*^(a), 1973).

Conclusion

The acute oral toxicity of benzoyl peroxide is very low: LD_{50} in mice >2,000 mg/kg, and in rats 5,000 mg/kg. No deaths occurred in male rats following inhalation of 24.3 mg/l.

3.1.3 Irritation

<u>Skin Irritation</u>

In a rabbit study in which 78 % granular benzoyl peroxide was applied to the back of 3 males and 3 females, the substance was held in place for 4 hours with a gauze bandage. After 4 hours, the bandages were removed and the exposed areas washed with lukewarm water. The skin was examined for any injury or irritation according to Draize scale at 4, 24 and 72 hours. No signs of skin irritation were observed at 1, 3 days after 4 hr exposure (Wazeter, F.X., et al(c), 1973). But 24 hrs-patch tests of different cases indicated that benzoyl peroxide has an irritant potential. Pure benzoyl peroxide and 10 % solution of benzoyl peroxide in propylene glycol led to slight erythema in guinea pigs under patches for 24 hrs (Eastman Kodak Co., 1964). In another 24 hrs-patch test, different concentrations of benzoyl peroxide in yellow soft paraffin were applied to shaved areas on the back of rabbits under occlusion for 24 hrs to determine which causes an erythematous reaction after single epicutaneous application in 50 % of the treated animals. The lowest benzoyl peroxide irritant concentration was 1 % for one animal. All rabbits reacted with 10 %, 15 % and 30% concentrations without exception. The ID50 of benzoyl peroxide was 2.52 %. This result suggests that benzovl peroxide is a skin irritant (Haustein, U.F., et al., 1985). In a cumulative skin irritation study, when 100 µl of 10 % Benzoyl peroxide was applied to the entire dorsum of mice for 6 months, keratinocytes showed marked variations in size and shape (atypia). Some were degenerating and separating each other.

A focal serous crust lay over the epidermis (Kligman, A.M., et al, 1998). Another 21-day cumulative test to determine the skin irritation profiles of benzoyl peroxide when entrapped in the polystyrene system and dispersed in oil-in-water vhicles was conducted. The cumulative 21-day irritancy score in six rabbits per group showed reduced skin irritation with the entrapped/controlled-release system (Wester RC et al, 1991).

Eye Irritation

In a study on eye irritation in rabbits, 70 % granular benzoyl peroxide was not irritating to the eye of 3 rabbits when immediately washed after instillation, whereas one rabbit in which benzoyl peroxide was not washed out immediately showed eye irritation until 72 48 hrs after application of 100 mg benzoyl peroxide. Corrosiveness was not observed in all animals (NIER, Korea, $2002^{(f)}$). In another eye irritation test, 8 male and female rabbits were divided into 2 groups. The 5 rabbits in group 1 were exposed to 111.4 mg benzoyl peroxide for 5 minutes. The rabbits in group 2 were exposed to 120.7 mg of the substance for 24 hrs. The study concluded that benzoyl peroxide was not irritating to the eyes of albino rabbits if it was washed out within 5 minutes after being placed in the conjunctival sac, however, if 78 % benzoyl peroxide was not washed out until 24 hr later, it proved to be a strongly irritating (Wazeter, F.X., *et al*^(d), 1973).

Conclusion

Benzoyl peroxide was slightly irritating to skins in 24 hr-patch tests. Benzoyl peroxide was not irritating to the eyes of rabbits if washed out within 5 minutes after instillation, however, if the chemical was not washed out until 24 hours later, it proved to be irritating.

3.1.4 Sensitisation

Studies in Animals

Several skin sensitization tests have been performed with benzoyl peroxide according to standard protocols, with a good concordance (table 4). In a study with mouse, benzoyl peroxide was evaluated in each of five independent laboratories using either the standard murine local lymph node assay (LLNA) protocol (Kimber, I., *et al*, 1998 & Basketter, D.A, *et al*, 1996) or minor modifications of it. Stimulation indices (SI) of 3 (EC₃ values) or greater was considered to have skin sensitizing activity. Benzoyl peroxide elicited strong LLNA response in all trials by 5 laboratories. Even at the lowest concentration tested (0.5 %), benzoyl peroxide induced stimulation indices of considerably greater than 3, and for this reason it proved to be impossible to derive EC₃ values mathematically (Kimber, I., *et al*, 1998).

Table 4. Skin sensitization s	studies on bezoyl	peroxide
-------------------------------	-------------------	----------

Methods	Species	Results	References
LLNA	Mouse	Positive	Kimber, I., et al., 1998
Buehler test	Guinea pig	Positive	Basketter, D.A, et al., 1996
TINA test	Guinea pig	Positive	Haustein, U.F., et al., 1985
Maximization test	Human	Positive	Leyden, J.J., et al., 1977

<u>Studies in Humans</u>

Allergic contact dermatitis from benzoyl peroxide in a therapeutic as well as an occupational context is well documented. Several positive results were obtained in the patch tests performed on human patients with leg ulcers or acne (Eaglstein, W.H., 1968, Von Bahmer, F.A., *et al*, 1984, Morillas, I.M., *et al*, 1987, Agathos, M., *et al*, 1984, Rietschel, R.L. *et al*, 1982). Loosening of a hip prothesis was reported from allergy to benzoyl peroxide used as an acivator of acrylic bone cement (Jäger, M., *et al*, 1979, Mosby Year Book, 1992). A few cases of allergic reaction were reported in dentists, bakers or other workers who might be exposed to benzoyl peroxide in their working environment (Bonnekoh, B., *et al*, 1984, Santiago, Q., *et al*, 1993, Gebhardt, M., *et al*, 1996)

In a human maximization test, 4 formulations containing benzoyl peroxide, two 5 % and two 10 % gel preparations for acne, were assayed for contact sensitizing capability (Leyden, J.J, *et al*, 1977). 38 of 50 subjects became sensitized and the sensitized subjects reacted to all four formulations. The authors, meanwhile, referred that the actual rate of seinsitization in clinical usage is dependent on many variables such as rapid loss and dilution by facial sweat, instability of benzoyl peroxide, which all minimize the likelihood of sensitization.

3.1.5 Repeated Dose Toxicity

A study conducted by NIER was identified as the key study because it was well conducted under the GLP using OECD test guideline 422 (NIER^(d), Korea, 2001). In this study, the rats were exposed to benzoyl peroxide at levels of 0, 250, 500 and 1,000 mg/kg/day for 29 days for males and for 41-51 days for females. No deaths were found in any animals including control groups during exposure period. Hematemesis was observed in one male receiving 500 mg/kg but this is not related to benzoyl peroxide exposure. No hematological effect attributable to benzoyl peroxide was observed in any treated groups. A significant dose-related decrease in the weight of testes and epididymis wasobserved in males treated with 1,000 mg/kg. In the males, receiving 1,000 mg/kg, testis degenerations were observed microscopically(5 animals of 10). In females at 1,000 mg/kg, slight effects in uterus such as epithelial vacuolation or hyperplasia were observed(control 0/10; 250 mg/kg 0/10, 500 mg/kg 2/10; 1000 mg/kg 3/10). There were no significant dose-related changes in any other pathological findings. Since reproductive organ weights were significantly changed in treated animals this study suggested the NOAEL as to 500 mg/kg for male rats.

In a long term feeding study (Sharratt, M., *et al*, 1964), groups of rats and mice were fed on diets of wholemeal flour and breadcrumbs made with treated flour containing benzoyl peroxide. The flour contained 2800 ppm, 280 ppm, 28 ppm and 2.8 ppm of benzoyl peroxide. The experiment was terminated after 80 and 120 weeks treatment in mice and rats, respectively. All animals that were killed at these times, or that died were autopsied. No effect on weight gain was found during the first 12 months in rats eating bread prepared from flour treated with benzoyl peroxide. The reduced rate of weight gain was seen in the rats whose diets contained flour treated with 2,800 ppm and 280 ppm benzoyl peroxide. The study indicated that the flour treated with 28 ppm benzoyl peroxide had no significant effect on the rate of weight gain of rats. There was no significant difference between treated and control groups in the mortality of the animals(rat and mice) except the group receiving 280 ppm benzoyl peroxide, in which there were a large number of accidental deaths. There were statistically significant incidence of atrophy of the testicles in the rats given the diet based on flour treated with benzoyl peroxide at 2800 ppm and 2.8 ppm. Accurate amount of dose cannot be determined because actual consumption amount of the diet was not described in the study.

Conclusion

Repeated administration by oral gavage to 1,000 mg/kg/day for 29 days resulted in decreased weights of testes and epididymis in male rats. The NOAEL for repeated dose toxicity is 500 mg/kg/day.

3.1.6 Mutagenicity

In vitro Studies

Several *in vitro* studies show that benzoyl peroxide is non-mutagenic to test strains *Salmonella typhimurium* TA97, TA98, TA100, TA102, TA104, TA1535, TA1537, and *Escherichia coli* WP2 uvrA in the presence and absence of a metabolic activation system (NIER^(e), Korea, 2001; Fujita, H., *et al.*, 1987; Dillon, D., *et al.*, 1998; Zeiger, E., *et al.*, 1988). Also one mammalian chromosomal aberration test *in vitro* with Chinese Hamster Lung (CHL) cells yields a negative result without metabolic activation at 0.2 mg/ml, the upper limit concentration to allow the cell proliferation (Ishidate, M.J., *et al.*, 1980).

In vivo Studies

An *in vivo* study, a mammalian erythrocytes micronucleus test, was conducted in accordance with OECD TG 474 and GLP. Benzoyl peroxide was administered once by intraperitoneal injection to male and female ICR mice at doses up to 200 mg/kg b.w. Benzoyl peroxide did not influence the incidence of micronuclei formation up to a maximum concentration of 200 mg/kg (NIER^(f), Korea, 2001) Similarly, a dominant lethal assay with the ICR mouse strain was also negative at 54 and 62 mg/kg (Epstein, J., *et al.*, 1972).

Conclusion

Benzoyl peroxide does not cause gene mutations in bacteria nor *in vitro* chromosomal aberrations in Chinese Hamster Lung (CHL) cells. In a mammalian erythrocytes micronucleus test performed *in vivo*, negative results were obtained. It can be concluded that benzoyl peroxide is not genotoxic.

3.1.7 Carcinogenicity

In vivo Studies in Animals

Dermal

A two-year dermal carcinogenicity study conducted according to GLP found no evidence of oncongenicity resulting from daily topical application of a benzoyl peroxide gel in mice at doses that meet the maximum tolerance dose(CHPA, 2001). In this study benzoyl peroxide was applied at doses of 1, 5 and 25 mg per mouse once daily for 104 weeks to a 2 x 3 cm area on the dorsal skin of animal. Discontinuous-treatment groups were applied the high dose of benzoyl peroxide, 25 mg per mouse, for 52 weeks and the vehicle for the rest of the study period. The animals were sacrificed at 52 weeks (interim sacrifice) or 104 weeks, and complete necropsies were performed. Benzoyl peroxide had no effect on survival, weights, food consumption, or gross pathology, except for treatment-site ulceration in the high- dose group, and produced no evidence of systemic toxicity. Microscopic evaluation revealed treatment-related findings confined to the site of application. The study concluded that there were no treatment related adverse effects in any of the neoplastic lesions analyzed in either sex.

Benzoyl peroxide was first reported to have the possible activity of promoter of cutaneous papilloma and squamous cell carcinomas in female SENCAR mice initiated with DMBA (Slaga, T.J., *et al.*, 1981) at the concentrations of 1, 10, 20 and 40 mg of benzoyl peroxide for 52 weeks. In another study, tumor-promoting activity was found in 30 SENCAR mice initiated with 10 nmols of DMBA with 0.1, 0.3, 0.6, 2.0 and 20 mg of benzoyl peroxide given twice a week for 100 weeks. From this study, the lowest tumor promoting dose was 0.6 mg, and no tumors were induced in initiated mice treated with either 0.1 or 0.3 mg of benzoyl peroxide (Gimenez-Conti, I.B., *et al.*, 1993). Including these *in vivo* mice skin tumor studies, relatively consistent positive results have been found in skin tumor promotion experiments in which SENCAR mice, one of the most sensitive types, were treated with experimental initiation agents such as 7,12-dimethyl benz[a]anthracene (DMBA) or N-methyl-N'-nitrosoguanidine (MNNG) (Slaga, T.J., *et al.*, 1981; Gimenez-Conti, I.B., *et al.*, 1993; Sharratt, M., *et al.*, 1964; O'Connell, J.F., *et al.*, 1986; Kurokawa, Y., *et al.*, 1984). These *in vivo* mice studies suggest that benzoyl peroxide may not be a complete carcinogen and/or initiator, but it seems to be a tumor promoter in skin of mice.

However, Kurokawa et al. observed significant increases in the incidence of skin tumors when benzoyl peroxide alone was applied at 20 mg for 51 weeks. The first skin tumor was observed in week 24 in a mouse treated with benzoyl peroxide. At the end of study, 8 out of 20 mice developed skin tumors related to the substance, of which 5 had squamous cell carcinomas. They concluded that the chemical has possible complete carcinogenic properties as well as a definite promoter activity(Kurokawa, Y., *et al.*, 1984). On the other hand, the published studies of potential interactions between benzoyl peroxide in gel and lotion vehicles, typical types of medications for acne, and UV radiation in mouse skin provide no evidence that benzoyl peroxide promoted or enhanced UV radiation-induced skin carcinogenesis (Iversen, O.H., *et al.*, 1988; Epstein, J., *et al.*, 1988).

Oral

Two oral carcinogenicity studies were also conducted by Sharratt*et al.*, (1964). In these studies, benzoyl peroxide was added to the diet in the form of commercial flour bleach at levels of 28, 280, or 2,800 mg/kg diet for 120 weeks (albino rats) or 80 weeks (albino mice) and there was no evidence of compound–related tumor induction. In the same *in vivo* study, single subcutaneous injection of benzoyl peroxide suspended in starch solution did not induce exposure-related tumors. These other exposure experiments also elicited the negative findings of carcinogenicity (Sharratt, M., *et al.*, 1964).

Studies in Humans

In the case of *in vitro* studies, benzoyl peroxide has been observed to induce DNA single strand breaks, which is an indication that treatments lead to events reducing the average size of single-strand DNA fragments in treated cells. For example, in normal human bronchial epithelial cells, the induction of DNA single strand breaks was observed at the concentration shown to inhibit growth (Saladino, A.J., *et al.*, 1985). One pilot case-control epidemiological study with 159 cases and 213 age- and sex-matched controls showed no significant association between the prescribed use of benzoyl peroxide to treat acne and malignant melanoma, or between the occurrence of acne and malignant melanoma (Cartwright, R.A., *et al.*, 1988). In a much larger case-control study, no statistically significant association was found between all prescription and nonprescription use of benzoyl peroxide-containing acne medications and skin cancer involving 964 skin cancer cases and 3856 controls (Hogan, D.J., *et al.*, 1991).

<u>Conclusion</u>

There is no evidence to suggest that benzoyl peroxide is a carcinogen. However, there is some evidence from non-guidelines studies that benzoyl peroxide is a skin tumour promoter.

3.1.8 Toxicity for Reproduction

A combined repeated dose and reproduction/developmental toxicity screening test (OECD TG 422) was conducted using Sprague-Dawley rats (NIER^(d), Korea, 2001). Benzoyl peroxide was administered to rats by gavage at doses of 0, 250, 500 and 1,000 mg/kg/day. Males were dosed for 29 days and females were dosed for 41 - 51 days from 14 days before mating to day 3 of lactation throughout the mating and pregnancy period. No treatment-related changes in precoital time and rate of copulation, fertility and gestation were noted in any benzoyl peroxide treated groups. Minimal symptoms such as vacuolation or hyperplasia, were seen in 1000 mg/kg group, but this is not considered to have been related to benzoyl peroxide treatment. Adverse effects on reproduction were shown at the highest dose of 1,000 mg/kg/day in male rats with the reduction of reproductive organ's weight and slight testes degenerations. The Sharratt et al study referred in section 3.1.3, indicating testicular atrophy in rates treated with benzoyl peroxide support the result of this study. In female rats, no adverse effects were observed during the test period. The NOAEL for reproduction toxicity in male rats was 500 mg/kg/day. No variants were found. High birthrate of runt was seen and body weight gain of pups was significantly decreased (male 9 %; female 12.9 % of control wt) at dose of 1,000 mg/kg/day. The study concluded that Benzovl peroxide has adverse effects on development of pups with high birthrate of runt at 1000 mg/kg dose level. The NOAEL for developmental toxicity was determined as 500 mg/kg/day.

Conclusion

The NOAEL for reproduction toxicity in male rats is 500 mg/kg based on reduced weight of testes and epididymis at 1,000 mg/kg. NOAEL for developmental toxicity is also 500 mg/kg based on significant decreased weight gain of pups with high birthrate of runt at 1,000 mg/kg.

3.2 Initial Assessment for Human Health

In Korea, total production volume of benzoyl peroxide was 1,357 tonnes in 2001, and the chemical is produced by only one company. The amounts of import and export were estimated as 268 and 293 tonnes/year, respectively. 75 % of benzoyl peroxide is mainly used as an initiator of polymerization in the manufacture of expandable styrene polymer and other resins. Benzoyl peroxide is used in the treatment of acne vulgaris and the medical products contain mainly 5 - 10 % benzoyl peroxide. A very small portion of benzoyl peroxide is used as a flour bleaching agent.

The acute oral toxicity of benzoyl peroxide is likely to be low since the LD_{50} is higher than 2,000 mg/kg in mice and 5,000 mg/kg in rats. Inhalation LC_0 is 24.3 mg/l in male rats with observable signs of eye squint, dyspnea, salivation, lacrimation, erythema and changes of respiratory rates and motor activity.

Benzoyl peroxide was slightly irritating to skin of test animals in 24 hour-patch tests. Benzoyl peroxide was not irritating to the eyes of rabbits if it was washed out within 5 minutes after instillation, however, if the chemical was not washed out until 24 hr later, it proved to be irritating.

Benzoyl peroxide was skin sensitizing in mice and guinea pigs when tested using the LLNA (Local Lymph Node Assay) and Buehler test or in human volunteers using the maximization test.

In the combined repeated dose and reproduction/developmental toxicity study (OECD TG 422), benzoyl peroxide did not lead to hematological- and biochemical-related adverse effects. Repeated administration by oral gavage to 1,000 mg/kg/day for 29 days resulted in decreased weights of testes and epididymis in male rats. The NOAEL for repeated dose toxicity was 500 mg/kg/day.

This substance did not cause gene mutation in bacteria (OECD TG 471 & 472) and *in vitro* chromosomal aberration in CHL (Chinese Hamster Lung) cells. An *in vivo* mammalian erythrocytes micronucleus test (OECD TG 474) showed negative results. This means that benzoyl peroxide is not genotoxic.

Benzoyl peroxide is not likely to be a complete carcinogen and/or initiator, but it seems to be a tumor promoter in mouse skin in an experimental two-stage model of carcinogenesis.

In the above combined repeated dose and reproduction/developmental toxicity study, no treatmentrelated changes in precoital time, rate of copulation, fertility and gestation were noted in any treated group. The adverse effects on reproduction were shown at the highest dose of 1,000 mg/kg/day in male rats with the reduction of reproductive organ's weight and slight testes degenerations. In female rats, no adverse effects were observed during the test period. The NOAEL for reproduction toxicity in male rats was 500 mg/kg/day. There was no evidence of teratogenic effects of benzoyl peroxide, but the body weight gain of pups at a dose of 1,000 mg/kg/day was significantly decreased. The NOAEL for developmental toxicity was 500 mg/kg/day.

The major routes of occupational exposure are inhalation and dermal. Limited data on exposure are available. Annual monitoring of the concentration of benzoyl peroxide at the workplace of the production facility has shown concentrations in airborne aerosols (personal sampling) of less than 1 mg/m^3 .

4 HAZARDS TO THE ENVIRONMENT

4.1 Aquatic Effects

Acute Toxicity Test Results

The acute toxicity data of benzoyl peroxide to fish, daphnia, algae and microorganisms were investigated for the aquatic effects. Only a few studies with the substance were available. Results of aquatic effects are summarized in Table 5.

Two acute Algae growth inhibition tests (OECD TG 201) showed comparable results. The 72 hr- E_bC_{50} (biombass) of *Selenastrum capricormutum* was 0.07 mg/L and the 72 hr- ErC_{50} (growth rate) was 0.44 mg/L in a static system based on nominal concentrations (NIER, Korea, 2001f). This test was well documented and performed according to OECD test guidelines and GLP . It can be considered to be valid without restrictions. In another study using *Pseudokirchneriella subcapitata*, a 72 hr- E_bC_{50} of 0.44 mg/L and a 72 hr- E_rC_{50} of 0.83 mg/L had been derived in static test (Akzo Nobel, the Netherlands).

Organism	Species	Test Duration	Result (mg/L)	Remarks	Reference
Almo	Selenastrum capricornutum	72 hr	$EbC_{50} = 0.07$ $ErC_{50} = 0.44$	OECD TG 201 (static, nominal concentration)	NIER, Korea, 2001f
Algae Pseudokirch ella subcapii		72 hr	$EbC_{50} = 0.44$ $ErC_{50} = 0.83$	OECD TG 201 (static, nominal concentration)	Akzo Nobel, The Netherlands
	Daphnia Magna	48 hr	$EC_{50} = 0.07$	OECD TG 202 (static, nominal concentration)	NIER, Korea, 2001g
Invertebrates	Daphnia Magna	48 hr	EC ₅₀ = 2.91	OECD TG 202 and EEC Directives (static, nominal concentration)	Wijk, R.J. van <i>et al</i> , 1999
Fish	Oryzias latipes	96 hr	$LC_{50} = 0.24$	OECD TG 203 (flow-through, mean measured concentration)	NIER, Korea, 2002c
1 1511	Poecilia reticulata	96 hr	$LC_{50} = 2.0$	OECD TG 203 and EEC method, part C (semi-static, nominal concentration)	Mark, U <i>et al</i> , 1989
Microorganisms	activated sludge	30 min.	$EC_{50} = 35$	OECD TG 209	Ginkel, C.G. van <i>et al</i> , 1990

 Table 5. Effects of Benzoyl peroxide on Aquatic Organisms

The 48 hr-EC₅₀ of *Daphnia magna* in an immobilization test (OECD TG 202) was calculated to be 0.07 mg/L. The test was performed in a static system based on nominal concentrations (NIER,

Korea, 2001g). A 48 hr-EC₅₀ of 2.91 mg/L has been derived for *Daphnia magna* in another test with a static system (Wijk, R.J. van *et al*, 1999).

A 96 hr-LC₅₀ of 2.0 mg/L has been derived for guppies (*Poecilic reticulata*) in a semi-static test based on nominal concentration of 0.7, 1.3, 2.4 and 4.2 mg/L (Mark, U *et al*, 1989). In a study with *Oryzias latipes*, the 96 hr-LC₅₀ was calculated to be 0.24 mg/L. The nominal concentrations were 0.25, 0.5, 1.0, 2.0 and 4.0 mg/L but the mean measured concentrations of benzoyl peroxide in the test solution of the flow-through system were 0.23, 0.47, 0.69, 1.54, and 2.17 mg/L, respectively (NIER, Korea, 2002c). In another study, a 96 hr-LC₅₀ of 3.9 mg/L was reported using a static system based on nominal concentration (MOE, Korea, 1996). These results indicated that different test results could be obtained according to the test protocols because benzoyl peroxide is very unstable in water (half-life : 5.20 hr at pH 7.0).

From the chemical structure, benzoyl peroxide is expected to be hydrolyzed to benzoic acid. The toxicity of benzoic acid to aquatic organisms is as follows (see corresponding SIAR for benzoic acid ,CAS No. 65-85-0);

Acute toxicity to fish: LC_{50} (96 hr) for *Lepomis macrochirus* is 44.6 mg/L and for *Salmo gairdneri* is 47.3 mg/L in static conditions.

Acute toxicity to aquatic invertbrates: EC_{50} (48 hr) for *Daphnia magna* is 100 mg/L in static conditions.

Acute toxicity to aquatic plants: EC_{50} (3 h) for *Scenedesmus quadricauda* is 75 mg/L (inhibition of photosynthesis).

Conclusion

Benzoyl peroxide is toxic to aquatic organisms. On the acute toxicity basis, the most sensitive aquatic organism is algae with an E_bC_{50} in the range of 0.07 - 0.44 mg/L.

4.2 Terrestrial Effects

Toxicity to terrestrial organisms was tested in a limited number of species. An earthworm acute toxicity test (OECD TG 207) was performed with *Eisenia foetida* on artificial soil. The 14d-LC₅₀ for *Eisenia foetida* was more than 1,000 mg/kg because the limit test of 1,000 mg/kg showed no mortality (NIER, Korea, 2002d). This toxicity value suggests that toxicity of benzoyl peroxide to soil dwelling organisms would be low.

Toxicity to non-mammalian terrestrial species (Chicken embryo) was investigated. For the toxic effect upon embryos, benzoyl peroxide was injected into the shell membrane of 3-day chicken eggs in the air chamber for 14 days. The 2d-LD₅₀ for chicken embryos was 0.99 μ moles/egg and the 14d-LD₅₀ was 0.82 μ moles/egg (Korhonen, A, *et al*, 1984).

Information on effects of benzoyl peroxide on birds, mammals, plants, invertebrates and other terrestrial organisms are not available.

4.3 Other Environmental Effects

No data available

4.4 Initial Assessment for the Environment

Benzoyl peroxide is commercially produced as a white granule with purities ranging from 22 to 99 %. It has a melting point of 104 - 106 °C, vapour pressure of 0.00929 Pa, solubility of 9.1 mg/L in water at 25 °C, and log P_{ow} of 3.43 at 25 °C. In indirect photolysis, half-life is estimated to be 3 days by the AOPWIN model. The substance is readily biodegradable (OECD TG 301C: 83 % as BOD after 21 days) and hydrolyses rapidly in water [OECD TG 111] with a half-life of 11.87 hrs at pH 4.0 and 5.20 hrs at pH 7.0 at 25 °C. The main hydrolysis product of benzoyl peroxide is benzoic acid (a SIDS assessment of benzoic acid is available, CAS No 65-85-0). An estimated BCF of 92 with the BCFWIN model suggests that this chemical has a low potential for bioaccumulation. A generic fugacity model (Mackay level III) was used for environmental fate estimation. If the most realistic emission pattern to water is assumed then the substance will remain in the aquatic compartment.

Benzoyl peroxide has toxic effects on the aquatic organisms. The range of 72 hr- E_bC_{50} (biomass) and 72 hr- ErC_{50} (growth rate) values for algae is 0.07 - 0.44 mg/L and 0.44 - 0.88 mg/L. The 48 hr- EC_{50} for daphnia is 0.07 - 2.91 mg/L and the LC_{50} for acute toxicity to fish is 0.24 - 2.0 mg/L. Although benzoyl peroxide is highly toxic to aquatic organisms, the substance is not bioaccumulated because of the rapid removal by hydrolysis (half-life = 5.2 hr at pH 7 at 25 °C) and biodegradation. The toxicity observed is assumed to be due to benzoyl peroxide rather than benzoic acid, which shows much lower toxicity to aquatic organisms. One can assume that effects occur before hydrolysis takes place. The toxicity for terrestrial organisms (earthworms) is low (14d- $LC_{50} > 1,000$ mg/kg for *Eisenia foetida*).

5 **RECOMMENDATIONS**

<u>Human Health</u>: The chemical possesses properties indicating hazards to human health (sensitisation, effect on testes weight, fetal body weight and skin tumour promotion activity) and is a candidate for further work, i.e. exposure assessment, and if considered necessary, risk assessment.

<u>Environment</u>: It is expected that the possibility of any environmental releases of benzoyl peroxide is low in the sponsor country. However, this substance is a candidate for further work, even if it hydrolyses rapidly and has a low bioaccumulation potential. The substance shows high acute toxicity to aquatic organisms and some information indicates wide-dispersive use of this substance. This could lead to local concern for the aquatic environment and therefore environmental exposure assessment is recommended.

6 REFERENCES

Agathos, M., Bandmann, H.J., Benzoyl peroxide contact allergy in leg ulcer patients, Contact Dermatitis, 17, 316 - 317, 1984

Akzo Nobel, The Netherlands, Effects of the Water-Accomodated Fraction of Lucidol on the Growth of the Freshwater Green Algae *Pseudokirchneriella subcapitata*, 1989

Aldrich, Catalog handbook of fine chemicals, Milwaukee, WI, Aldrich chem. Co., 163, 2000 - 2001

Autoxidized Squalene and Linoleic Acid, and of Simpler peroxides, in Relation to Toxicity of Radiation, Biochem. J. 67, 551 - 558, 1957

Basketter, D.A, Gerberick, G.F., An interlaboratory evaluation of the Buehler test for the identification and classification of skin sensitizers, Contact Dermatitis, 35, 146 - 151, 1996

Bonnekoh, B. and Merk, H.F., Airborne allergic contact dermatitis from benzoyl peroxide as a bleaching agent of candle wax, Contact Dermatitis, 24, 367 - 368, 1984

Cartwright, R.A., Hughes, B.R. and Cunliffe, W.J., Malignant Melanoma, Benzoyl Peroxide and Acne : A Pilot Epidemiological Case-Control Investigation, British J, Derm., 118, 239 - 242, 1988

Chem. Inspect Test Inst ; Biodegradation and Bioaccumulation Data of Existing Chemicals Based on the CSCL Japan ; Published by Japan Chemical Industry Ecology-Toxicology & Information Center., ISBN 4-89074-101-1, 1992

Dillon, D., Combes, R. and Zeiger, E., The Effectiveness of *Salmonella* strains TA100, TA102 and TA104 for Detecting Mutagenicity of Some Aldehydes and Peroxide, Mutagenesis, 13 (1), 19 - 29, 1998

Eaglstein, W.H., Allergic Contact Dermatitis to Benzoyl Peroxide, Arch Derm., 97, p 527, 1968

Epstein, J., Photocarcinogenesis Promotion Studies with Benzoyl Peroxide and Croton Oil, J. Investigative Dermatology, 91 (2), 114 - 116, 1988

Epstein, S.S., Arnold, E., Andrea, J., Bass, W. and Bishop, Y., Detection of Chemical Mutagens by the Dominant Lethal Assay in the Mouse, Toxicology and Applied Pharmacology, 23, 288 - 325, 1972

Fujita, H. & Sasaki, M., Mutagenicity Test of Food Additives with *Salmonella typhimurium* TA 97 and TA102(II), Annu. Rep. Tokyo Metro. Res. Lab. P.H., 38, 423 - 430, 1987 (in Japanese with English abstract)

Gebhardt, M., Geier, J., Evaluation of patch test results with denture material series, Contact Dermatitis, 34, 191 - 195, 1996

Gigiena, Sanitariya. V/O Mezhdunarodnaya Kniga 113095 Moscow, USSR Vol.1-, 1936 cited in SAX'S Dangerous Properties of Industrial Materials, Tenth Edition, 1999 John Wiley & Sons, Inc CD version

Gimenez-Conti, I.B., Oubari, D., and Slaga, T.J., Dose-response of benzoyl peroxide hyperplastic promoter and possible complete carcinogenic activities, Proc. Am. Assoc., Cancer Res., 34, 184, 1993

Ginkel, C.G. van and Stroo, C.A., Akzo Chemicals BV, The Netherlands, Toxicity of Dibenzoyl peroxide for activated sludge, 1990

Haustein, U.F., Kramer, I., Fickweiler, E., Beahandlung der Akne vulgaris mittels Benzoylperoxid, Dermatol. Monatsschr., 167, 149 - 153, 1981

Haustein, U-F., Tegetmeyer, L. and Ziegler, V., Allergic and Irritant Potential of Benzoyl Peroxide, Contact Dermatitis 13, 252 - 257, 1985

Horgan, V.J., Philpot, J. ST L., Porter, B.W. and Roodyn, D.B., Toxicity of Autoxidized Squalene and Linoleic Acid, and of Simpler peroxides, in Relation to Toxicity of Radiation, Biochem. J., 67, 551-558, 1957

Hogan, D., Wilson, E., Miller, A., Benzoyl peroxide and facial skin cancer, Am. J. Epidemiol., 134, 772, Criteria for Recommended standard, Occupational Exposure to Benzoyl Peroxide, NIOSH, 1977

Hogan, D.J., To T, Wilson, E.R., Miller, A.B., Robson, D., Holfeld, K., Lane, P., A study of acne treatments as risk factors for skin cancer of the head and neck., Brit. J. Dermatol., 125, 343 - 348, 1991

Ishidate, M.J., Yoshikawa, K. and Soluni, T., Mutagenicity Testing of Food Additives-First Screening Data Hegen-to-dokusei, Mutagens Toxicol., 3, 82 - 90, 1980

Iversen, O.H., Skin Tumorigenesis and Carcinogenesis Studies with 7,12dimethylbenzene[a]anthracene, ultraviolet Light, Benzoyl Peroxide (Panoxy gel 5 %) and Ointment Gel Carcinogenesis 9 (5), 803 - 809, 1988

Jager, M., and Balda, B.R., Loosening of a Total Hip Prosthesis at Contact Allergy Due to Benzoyl Peroxide, Arch. Orthop. Traumat. Surg., 94, 175 - 178, 1979

Kimber, I., Hilton, J., Dearman, R.J., Gerberick, G.F., Ryan, C.A., Basketter, D.A., Lea, L., House, R.V., Ladics, G.S., Loveless, S.E. and Hastings, K.L., Assessment of the Skin Sensitization Potential of Topical Medicaments Using the Local Lymph Node Assay, An Interlaboratory Evaluation, J. Toxicol. Environ. Hlth., Part A, 53 (7), 563 - 579, 1998

Kligman, A.M. and Kligman, L.H., A Hairless Mouse Model for Assessing the Chronic Toxicity of Topically Applied Chemicals, Food and Chemical Toxicology, 36, 867 - 878, 1998

Korhonen, A., Hemminki, K. and Vainio, H., Embryotoxic Effects of Eight Organic Peroxides and Hydrogen Peroxide on Three-Day Chicken Embryos, Environ. Res., 33, 54 - 61, 1984

Kuchle, H.J., Untersuchungen der die augenschadigende Wirkung organischer Peroxyde, Zentralblatt fur Arbeitsmedizin und Arbeitsschuntz, Heit, 2, 25 - 31, 1958

Kurokawa, Y., Takamura, N., Matsushima, Y., Imazawa, T. and Hayashi, Y., Studies on the Promoting and Complete Carcinogenic Activities of Some Oxidizing Chemicals in Skin Carcinogenesis, Cancer Letters, 24, 299 - 304, 1984

Lewis, R.J Sr., Eds. Hawley's Condensed Chemical Dictionary, 14th Ed John Wiley & Sons, Inc., New York, 127 - 128, 2001

Leyden, J.J., Kligman, A.M., Contact sensitisation to benzoyl peroxide, Contact Dermatitis, Vol 3, 273 - 275, 1977

Lyman, W.J. *et al.*, Handbook of Chemical Property Estimation Methods. Washington DC, Amer. Chem. Soc., 5 - 4, 5 - 10, 1990

Marhold, J.V., Sbornik Vysledku Toxixologicheho Vysetreni Latek A Pripravku, Institut Pro Vychovu Vedoucicn Pracovniku Chemickeho Prumyclu Praha, Czechoslovakia, 1972, cited in SAX'S Dangerous Properties of Industrial Materials, Tenth Edition, 1999, John Wiley & Sons, Inc CD version

Mark, U., Meuwsen, I.J.B., Akzo Chemicals BV, The Netherlands, Acute Toxicity of Dibenzoyl peroxide to fish, 1989

Meylan WM, Howard PH, Environ Toxicol Chem 10, 1283 - 93, 1991

Morillas, I.M., Martinesz, A.A., Lozano J.L.S. and Perez A.G., Is benzoyl peroxide an irritant or sensitizer?, Contact Dermatitis, 16, 232 - 233,1987

Morsches, B. and Holzmann, H., Untersuchungen zur perkutanen Resorption von Benzoylperoxid, Arzneim.-Forsch./Drug Res., 32 (1), Nr.3, 1982

Mosby Year Book, Contact and Occupational Dermatology, St. Louis, MO/Marks, J.G., V.A. DeLeo, 164 - 164, 1992

Nacht, S., Yeung, D., Joseph, N., Beasley, J. N. Jr., Anjo, M. D. and Maibach, H.I., Benzoyl Peroxide : Percutaneous Penetration and Metabolic Disposition, J. Am. Acad. Dermatol. 4 (1), 31 - 37, 1981

National Institute of Environmental Research (NIER), Korea (2001a), "Hydrolysis as a Function of pH" tested by LGCI

National Institute of Environmental Research (NIER), Korea (2002a), Estimation of Physical Chemical Properties and Environmental Fate of SIDS Chemicals (II) - on Benzoyl peroxide

National Institute of Environmental Research (NIER), Korea (2001b), Acute oral toxicity of benzoyl peroxide, Report No. G01083 tested by KRICT

National Institute of Environmental Research (NIER), Korea (2001c), Combined Repeated Dose Toxicity with the Reproduction /Developmental Toxicity Screening Testing of Benzoyl peroxide in Rats, Report No.P049 tested by LGCI, 2001^(d)

National Institute of Environmental Research (NIER), Korea (2001d), Korea, Bacterial Reverse Mutation Test, Report No. S350 tested by LGCI

National Institute of Environmental Research (NIER), Korea (2001e), Micronucleus test of Benzoyl peroxide in mouse, Report No.S349 tested by LGCI,

National Institute of Environmental Research (NIER), Korea (2002b), Acute Eye Irritation/Corrosion Test in Rabbits, Report No. S367 tested by LGCI

National Institute of Environmental Research (NIER), Korea (2001f), The toxicity of Benzoyl peroxide to Aquatic plants (algae), Report No. EG01005 tested by KRICT, 2001^(h)

National Institute of Environmental Research (NIER), Korea (2001g), The acute toxicity of Benzoyl peroxide to Aquatic Invertebrate (Daphnia), Report No. EG01006 tested by KRICT, 2001⁽ⁱ⁾

National Institute of Environmental Research (NIER), Korea (2002c), The acute toxicity of Benzoyl peroxide to fish, Report No. EG01077 tested by KRICT

National Institute of Environmental Research (NIER), Korea (2002d), The toxicity of Benzoyl peroxide to earthworm tested by KRICT

NIOSH, Criteria for Recommended Standard, Occupational Exposure to Benzoyl Peroxide, 1977

O'Connell, J.F., Klein-Szanto, A.J.P., DiGiovanni, D.M., Fries, J.W. and Slaga, T.J., Enhanced Malignant Progression of Mouse Skin Tumors by the Free-Radical Generator Benzoyl Peroxide, Cancer Research, 46 (6), 2863 - 2865, 1986

OECD, Environment Directorate, Joint Meeting of the Chemicals Committee and the Working Party on Chemicals, Pesticides and Biotechnology, OECD Series on Testing and Assessment Number 27, Guidance Document on the Use of the Harmonised System for the Classification of Chemicals Which are Hazardous for the Aquatic Environment, ENV/JM./MONO(2001)8

Online Toxicology Data Network (TOXNET) : Hazardous Substances Data Bank (HSDB), 2002

Philpot, J. St. L. and Roodyn, D.B., A Comparison Between the Effects in Mice of Injected Organic Peroxides and of Whole-body X-Irradiation, Int. J. Rad. Biol., 4, 372 - 382, 1959

Rietschel, R.L. and Duncan, S.H., Benzoyl peroxide reactions in an acne study group, Contact Dermatitis, 8, 323 - 326, 1982

Saladino, A.J., Willey, J.C., Lechner, J.F., Grafstrom, R.C., LaVeck, M. and Harris, C.C., Effects of Formaldehyde, Acetaldehyde, Benzoyl Peroxide, and Hydrogen Peroxide on Cultured Normal Human Bronchial Epithelial Cells, Cancer Research, 45, 2522 - 2526, 1985

Santiago, Q., J.M. Olaguibel, B.E. Garcia and Tabar, A.I., Occupational airborne contact dermatitis due to benzoyl peroxide, Contact Dermatitis, 29, 165 - 166, 1993

Sevila, A., Romaguera C., Vilaplana J., Botella R., Contact dermatitis in children, Contact Dermatitis, 30, 292 - 294, 1994

Sharratt, M., Frazer, A.C. and Forbes, O.C., Study of the Biological Effects of Benzoyl Peroxide, Food Cosmet. Toxicol.2, 527 - 538, 1964

Slaga, T.J., Klein-Szanto, A.J.P., Triplett, L.L.I., Yotti, L.P. and Trosko, J.E., Skin Tumor-Promoting Activity of Benzoyl Peroxide, a Widely Used Free Radical-Generating Compound, Science 213, 1023 - 1025, 1981

Takahashi, A., Problems of Hygiene Maintenance for Food Coming into Contact with Rubber and Plastic Products, Nippon Gomu Kyokaishi 48 (9), 537, 1975 [Translated by English, E. A., Internat'l Polymer Sci. Technol. 3 (1), P 93 -105, 1976

The OECD HPV program, SIDS Initial Assessment Report for 13th SIAM- Benzoates category

Tkach, J,R., Allergic Contact Urticaria to Benzoyl Peroxide, Cutis 29, 187 - 188, 1982

Von Bahmer, F.A., Schulze-Dirks A. und Zaun H., Sensibilisierende Wirkung einer fur die behandlung des Ulcus cruris verwendeten 20 % igen Benzoylperoxid-Zubereitung, Dermatosen., 32, Nr 1, 21 - 24, 1984

Wazeter, F.X., and Goldenthal, E., Acute Inhalation to Rats, 328-005 International Research and Development Corporation (IRDC) August 31, 1973

Wazeter, F.X. and Goldenthal, E., Eye irritation in Albino Rabbits, International Research and Development Corporation (IRDC) August 31, 1973

Wazeter, F.X. and Goldenthal, E., Primary Skin Irritation and Corrosive Hazard Test in Albino Rabbits, International Research and Development Corporation (IRDC) August 31, 1973

Wazeter, F.X. and Goldenthal, E., Acute Oral Toxicity Test in Male Albino Rats, International Research and Development Corporation (IRDC) August 31, 1973

Weinberg, J.M., Moss, T., Gupta, S.M., White, S.M. and Don, P.C., Reticulate Hyperpigmentation of the Skin After Topical Application of Benzoyl Peroxide, Acta. Derm. Venereol (Stockh)., 78, 301 - 302, 1998

Wepierre, J., Corroller, M. and Didry, J. R., Distribution and Dissociation of Benzoyl Peroxide in Cutaneous Tissue after Application on Skin in the Hairless Rat, International Journal of Cosmetic Science 8, 97 - 104, 1986

Wester, R.C., Patel, R., Nacht, S., Leyden, j., Melendres, J. and Maibach, H., Controlled Release of Benzoyl Peroxide from a Porous Microsphere Polymeric System Can Reduce Topical Irritancy, J. Am. Acad. Dermatol., 24, 720 - 726, 1991

Wijk, R.J. van and Groenenboom, Ir. C. J., Akzo Nobel, The Netherlands, Acute Toxicity of Lucidol to Daphnia magna, 1999

Yeung, D., Nacht, S., Bucks, D. and Maibach, H. I., Benzoyl Peroxide: Percutaneous Penetration and Metabolic Disposition. II. Effect of Concentration, J. Am. Acad. Dermatol., 9, 920 - 924, 1983

Zeiger, E., Anderson, B., Haworth, S., Lawlor, T., Mortelmans, K. Salmonella Mutagenicity Tests: IV. Results from the Testing of 300 Chemicals. Env. Mol. Mutag., 11/12, 1 - 158, 1988

SIDS DOSSIER

Benzoyl peroxide

CAS No.: 94-36-0

Sponsor Country : Republic of Korea

Date of submission to OECD : August 2002

1.01 SUBSTANCE INFORMATION

*A.	CAS number	:	94-36-0
B.	Name (Primary	:	Benzoyl peroxide
nar	ne)		
*С.	Name (OECD	:	Benzoyl peroxide
nar	ne)		
† D.	CAS Descriptor	:	
E.	EINECS-Number	:	202-327-6
F.	Molecular Formula	:	$C_{14}H_{10}O_4$
*G.	Structural	:	
For	mula		



Smiles Code : O=C(OOC(=O)c(cccc1)c1)c(cccc2)c2

H. Substance Group

I. Substance Remark (Indicate the substance remark as prescribed in the EINECS Inventory, if possible)

J. Molecular Weight : 242.24

:

1.02 OECD INFORMATION

A.	Sponsor Country	:	Republic of Korea
B.	Lead	:	Environmental Risk Assessment Division, National Institute of
Org	anisation		Environmental Research
	Contact person	:	Dr. Hyun-Mi Kim
	Address		Environmental Research Complex
	Street	:	Gyeongseo-dong, Seo-gu
	Postal code	:	404-170
	Town	:	Incheon
	Country	:	Republic of Korea
	Tel	:	+82-32-560-7124
	Fax	:	+82-32-568-2037
	E-mail	:	hmikim@me.go.kr

C. Name of responder (Information on a responder should be provided when companies respond to Lead

Organisation or SIDS Contact Points.) Name : same as above Address : same as above

1.1 GENERAL SUBSTANCE INFORMATION

A.	Туре	of :	organic
1 1.	Lype	01 6	, organn

Substance

B. Physical State (at 20 °C and 1.013 hPa)

: solid

C. Purity	: 95 % (remainder water)
	75 % (remainder water)
	22 - 27 % (Mixture of 79 % benzoyl peroxide and some chemical
	additives for bleaching agent use)
Source	: Hansol chemical company (2001)

1.2 SYNONYMS

: Benzoic acid, peroxide Dibenzoyl peroxide Benzoperoxide Benzoyl Superoxide Benzoyl peroxide, remainder water Lucidol

1.3 IMPURITIES

No data

1.4 ADDITIVES

CAS No. EINECS No. Name Value Remarks	 7778-18-9 Calcium sulfate < 78 % Additive in 22 % benzoyl peroxide used in bleaching agent.
CAS No. EINECS No. Name Value Remarks	 471-34-1 Calcium carbonate < 78 % Additive in 22 % benzoyl peroxide used in bleaching agent.
CAS No. EINECS No. Name Value Remarks	 Starch < 78 % Additive in 22 % benzoyl peroxide used in bleaching agent.

1.5 QUANTITY

Estimated production	
	: Production in Korea : 1,375 tonnes/year
Remarks	: Imported into Korea : 268 tonnes/year
	Exported to EU: 293 tonnes/year
References	: National Institute of Environmental Research (NIER), Korea,
	survey on circulation volume and use pattern of benzoyl peroxide
	in Korea, 2001

1.6 LABELLING AND CLASSIFICATION

The following classifications have been established EEC Classification, Packaging, and Labeling of Dangerous Substances Official Journal of the European Communities, No L 136 (08 Jun. 2000)

EC No.	: 202-327-6
Index No.	: 617-008-00-0
Classification	: E, R2; Xi, R 36; R 43
Danger Symbol	: E, Xi
R-phrases	• R : 2-36-43
S-phrases	S : (2-) 3/7-14-36/37/39
Remarks	: Benzoyl peroxide is presently not classified as toxic chemical under the Toxic Chemical Control Act in Republic of Korea. It is in the process of being re- classified to toxic chemical because of its high aquatic toxicity.

1.7 USE PATTERN

А.	General	
	Type of Use	: Category
(a)	main	: non-dispersive use
	industrial	Plastic and polymer industry
	use	Initiator of polymerization
(b)	main	non-dispersive use
	industrial	Fabric industry
	use	Manufacture of sizing agents
(c)	main	: non-dispersive use
	industrial	Paints producing industry
	use	Manufacture of acrylic resin
(d)	main	: wide dispersive use
	industrial	Food industry
	use	Bleaching agents in flour
(e)	main	: wide dispersive use
	industrial use	pharmaceutical industry

OECD SIDS	BENZOYL PEROXIDE
1. GENERAL INI	ORMATION ID 94-36-0
	DATE: AUGUST 2002
	acne vularis treatment
Remarks	:
References	: National Institute of Environmental Research (NIER), Korea, survey on circulation volume and use pattern of benzoyl peroxide in Korea, 2001

B. Uses in Consumer Products

Function	Amount p	present	Physical state			
Bactericidal agent	2.5 - 10 %	%	gel type			
Remarks References	: National survey or in Korea,	Institute of n circulation v , 2001	Environmental volume and use	Research pattern of	(NIER), benzoyl p	Korea, peroxide

1.8 OCCUPATIONAL EXPOSURE LIMIT VALUE

Exposure limit value	
Туре	: TLV (Korea)
Value	: 5 mg/m^3 (8 hours TWA)
References	: Industrial safety and health act in Korea, 1998
Exposure limit value	
Туре	: TLV (US)
Value	: 5 mg/m^3 (8 hours TWA)
References	: ACGIH TLVs and Biological Exposure Indices for 1999
Exposure limit value	
Туре	: PEL
Value	: 5 mg/m^3
References	: OSHA Permissable Exposure Levels, CFR v.29 Section 1910.1000, 1996

1.9 SOURCES OF EXPOSURE

Source Remarks	 Manufacture Process : reaction (NaOH + H₂O₂ + 2C₆HCOCl) - Washing - Dewatering - Drying Possible release source : washing process (into wastewater treatment facility) Number of site : 1 Workplace exposure level : < 1 mg/m³ (as aerosol)
References	: Hansol Chemical Company in Korea

1.10 ADDITIONAL REMARKS

None

2.1 MELTING POINT

(a) Preferred Result	
Value :	= 104 - 106 °C Purity $= 97.\%$
Reliability	(2) Reliable with restrictions
References :	Aldrich, Catalog handbook of fine chemicals, Milwaukee, WI, Aldrich chem. Co., 164, 2000 ~ 2001
(b) Value	= 103 - 105 °C
Reliability	(2) Reliable with restrictions
References	Lewis, R.J. Sr., Eds., Hawley's Condensed Chemical Dictionary, 14th Ed, John Wiley & Sons, Inc. New York, 127 ~ 128, 2001
(c)	106.00
Reliability	(2) Reliable with restrictions
References	Richard, J., Lewis, Sr., SAX' Dangerous Properties of Industrial Materials. John Wiley & Sons, Inc. New York, on CD-ROM, 10th ed, 2000
(d) Value	- 105.90
value Reliability	$= 105 {}^{\circ}\text{C}$ (2) Reliable with restrictions
References	Lide, D.R. ed., CRC Handbook of Chemistry and Physics. 76th ed. Boca Raton, FL, CRC Press Inc., 3 ~ 250, 1995 ~ 1996
(e)	
Reliability	(2) Reliable with restrictions
References :	Budavari, S.ed., The Merck Index - An Encyclopedia of Chemicals, Drugs, and Biologicals, Whitehouse Station, NJ, Merck and Co., Inc., on CD-ROM version 12:2, 10th edition, 1998
(f) Value	-102 105 °C
Reliability	(2) Reliable with restrictions
References	Bretherick, L. Handbook of Reactive Chemical Hazards, Butterworths, 4th ed., London, 868 ~ 869, 1990
(g) Value	$-106 \cdot 108 \circ C$
Reliability	(2) Reliable with restrictions
References	Buckingham, J., ed., Dictionary of Organic Compounds, 5th ed., and Supplements, Chapman and Hall, New York, Vol. 2, 1612 ~ 1613, 1982

OECD SIDS	BENZOYL PEROX	IDE
2. PHYSICO-CHEMICAL D	ATA ID 94-3	36-0
	DATE: AUGUST 2	002
(h)		
Value	= 103 - 105 °C	
Reliability	: (2) Reliable with restrictions	
References	Bretherick, L., ed., Hazards in the Chemical Laboratory, 3rd	ed.
	The Royal Society of Chemistry, London, 270 ~ 271, 1981	
(i)		
Value	$= 103 ^{\circ}\mathrm{C}$	
Decomposition	: Yes	
Reliability	: (2) Reliable with restrictions	
References	U.S. Department of Health and Human Services and U	S.
	Department of Labor., NIOSH/OSHA Occupational Hea	ılth
	Guidelines for Chemical Hazards. 3 Vols., DHHS (NIOS	H)
	Publication No. 81 ~ 123, January, Volume 1, 1981	

2.2 BOILING POINT

References

(a) Preferred Result	
Value	: Decomposes explosively above 105 °C
Decomposition	: Yes
Reliability	: (2) Reliable with restrictions
References	: Lewis, R.J. Sr., ed., Hawley's Condensed Chemical Dictionary, 14th ed., John Wiley Sons, Inc., New York, 127 ~ 128, 2001
(b)	
Value	: Decomposes explosively
Decomposition	: Yes
Reliability	: (2) Reliable with restrictions

- : (2) Reliable with restrictions
- : Richard, J. Lewis, Sr. SAX' Dangerous Properties of Industrial Materials, on CD-ROM. 10th ed., John Wiley&Sons, Inc., New York, 2000

2.3 DENSITY (RELATIVE DENSITY)

(a)	
Туре	: Density
Value	: 1.33 g/cm^3
Temperature	: 25 °C
Reliability	: (2) Reliable with restrictions
References	: Lewis, R.J. Sr., ed., Hawley's Condensed Chemical Dictionary, 14th ed., John Wiley & Sons, Inc., New York, 127 ~ 128, 2001
(b)	
Туре	: Density
Value	: 1.33 g/cm^3
Temperature	: 25 °C

: (2) Reliable with restrictions Reliability

OECD SIDS	BENZOYL PEROXIDE
2. PHYSICO-CHEM	ICAL DATA ID 94-36-0
	DATE: AUGUST 2002
References	: International Agency for Research on Cancer, World Health
	Organization, IARC Monographs on the Evaluation of
	Carcinogenic Risk of Chemicals to Man.
	International Agency for Research on Cancer. Geneva. Vol. 36, 267 ~ 283, Supplement 7, p 58, 1999

2.4 VAPOUR PRESSURE

(a) Preferred Result Value Temperature Method Remarks Reliability References	 : = 0.00929 Pa : 25 °C : Estimated by MPBPWIN model : Input data (melting point = 105 °C) : (2) Reliable with restrictions : National Institute of Environmental Research (NIER), Korea, Estimation of Physical Chemical Properties and Environmental Fate of SIDS Chemicals (II)-on Benzoyl peroxide, 2002
(b) Value Temperature Reliability References	 Much less than 1 mmHg 20 °C (2) Reliable with restrictions U.S. Department of Health and Human Services and U.S. Department of Labor., OSHA Occupational Health Guidelines for Chemical Hazards, 3 Vols. DHHS (NIOSH) Publication No. 81 ~ 123, January, Volume 1, 1981

2.5 PARTITION COEFFICIENT LOG₁₀ P_{OW}

(a) Preferred Result	
Log P _{ow}	: = 3.43
Temperature	: 25 °C
Method	: Estimated by KOWWIN model
Remarks	: Input data (experimental Log $P_{ow} = 3.46$)
Reliability	: (2) Reliable with restrictions
References	: National Institute of Environmental Research (NIER), Korea, Estimation of Physical Chemical Properties and Environmental Fate of SIDS Chemicals (II)-on Benzoyl peroxide, 2002
(b)	
Log Pow	: = 3.46
Temperature	: 25 °C
Method	: other (measured) : no data
Reliability	: (2) Reliable with restrictions
References	: Online Toxicology Data Network (TOXNET), Hazardous

Substances Data Bank (HSDB), 2002

2.6 WATER SOLUBILITY

(a) Preferred Result	
Value : Temperature	9.1 mg/l 25 °C
Reliability :	(2) Reliable with restrictions
References :	 Online Toxicology Data Network (TOXNET), Hazardous Substances Data Bank (HSDB), 2002 Chem. Inspect Test Inst; Biodegradation and Bioaccumulation Data of Existing Chemicals Based on the CSCL Japan, Published by Japan Chemical Industry Ecology-Toxicology & Information Center, ISBN 4-89074-101-1, 1992
(b)	
Value :	Slightly soluble in water
Reliability:References:	(2) Reliable with restrictions Lewis, R.J. Sr., ed. Hawley's Condensed Chemical Dictionary, 14th ed., John Wiley & Sons, Inc. New York, 127 ~ 128, 2001
(c)	
Value :	Insoluble in water
Reliability :	(2) Reliable with restrictions Richard I Lewis Sr SAX' Dangerous Properties of Industrial
	Materials, on CD-ROM 10th ed., John Wiley & Sons, Inc., New York, 2000
(d)	
Value :	Sparingly soluble in water
Reliability : References :	(2) Reliable with restrictions Budavari, S. ed., The Merck Index - An Encyclopedia of Chemicals, Drugs and Biologicals on CD-ROM version 12:2, Whitehouse Station, NJ, Merck and Co., Inc., 10th edition, 1998
(e)	
Value :	Very sparingly soluble in water
Reliability :	(2) Reliable with restrictions International Agency for Research on Cancer World Health
	Organization. IARC Monographs on the Evaluation of Carcinogenic
	Risk of Chemicals to Man,
	International Agency for Research on Cancer. Geneva., Vol. 36, $267 \sim 283$, Supplement 7, p. 58, 1999
(f)	
Value :	Insoluble in water
References :	(2) Kenable with restrictions Bretherick L ed Hazards in the Chemical Laboratory 3rd ed. The
•	Royal Society of Chemistry, London, 270 ~ 271, 1981
2.7 FLASH POINT

(a)		
Value	: 40 °C	
Reliability	: (2) Reliable with restrictions	
References	: Baker, J., Material Safety Data Sheet (MSDS) Number, B1831 Effective Date, 11/02/01 by Mallinckrodt chemicals	
(b)		
Value	: 41 °C	
Reliability	: (2) Reliable with restrictions	
References	: Hazardous Chemical Database, hardy Research Group the University of AKRON, Department of Chemistry, 2001	
2.8 AUTO FLAMM	ABILITY	

(a)			
Value	: 80 °C		
Remarks	: 176 F		
Reliability	: (2) Reliable with restrictions		
References	: Lewis, R.J. Sr., ed., Hawley's Condensed Chemical Dictionary,		
	14th ed. John Wiley & Sons, Inc., New York, 127 ~ 128, 2001		
(b)			
Value	: 80 °C		
Remarks	: 176 F		
Reliability	: (2) Reliable with restrictions		
References	: Richard, J. Lewis, Sr. SAX' Dangerous Properties of Industrial Materials, on CD-ROM 10th ed., John Wiley & Sons, Inc., New York, 2000		
(c)			
Value	: 69 °C		
Remarks	: Calculated value		
Reliability	: (2) Reliable with restrictions		
References	: Bretherick, L. Handbook of Reactive Chemical Hazards, 4th ed.,		
	Butterworths, London, 868 ~ 869, 1990		

2.9 FLAMMABILITY

(a)	
Result	: Flammable
Remarks	: The decomposition product
Reliability	: (2) Reliable with restrictions
References	: Bretherick, L., Handbook of Reactive Chemical Hazards, 4th ed.,
	Butterworths, London, 868 ~ 886, 1990

OECD SIDS	BENZOYL PEROXIDE
2. PHYSICO-CHEMI	ID 94-36-0
	DATE: AUGUST 2002
(b) Posults	• Highly Flormable
Reliability References	 Fighty Flammable (2) Reliable with restrictions International Agency for Research on Cancer, World Health Organization. IARC Monographs on the Evaluation of Carcinogenic Risk of Chemicals to Man, International Agency for Research on Cancer, Geneva, Vol. 36, 267 ~ 283, Supplement 7, p. 58, 1999
(c) Results Remarks Reliability References	 Highly Flammable In the dry state (2) Reliable with restrictions National Fire Protection Association, Fire Protection Guide on Hazardous Materials, 7th ed., National Fire Protection Association Quincy, MA, 49 ~ 113, 49 ~114, 491M~57, 491M~145, 1981
(d) Results Remarks Reliability References	 Highly Flammable In the dry state (2) Reliable with restrictions U.S. Department of Health and Human Services and U.S. Department of Labor. NIOSH/OSHA Occupational Health Guidelines for Chemical Hazards, 3 Vol., DHHS (NIOSH) Publication No. 81 ~ 123, January, Volume 1, 1981

2.10 EXPLOSIVE PROPERTIES

(a)	
Results :	May explode spontaneously when dry (< 1 % of water)
Reliability :	(2) Reliable with restrictions
References :	Lewis, R.J. Sr., ed., Hawley's Condensed Chemical Dictionary, 14th ed., John Wiley & Sons, Inc., New York, 127 ~ 128, 2001
(b)	
Results :	May explode spontaneously when heated to above melting point, or when overheated under confinement. It is moderately sensitive to heat, shock, friction, or contact with combustible materials. Explosive decomposition above the mp forms flammable products.
Reliability :	(2) Reliable with restrictions
References :	Richard, J. Lewis, Sr. SAX' Dangerous Properties of Industrial Materials on CD-ROM 10th ed., John Wiley & Sons, Inc., New York, 2000
(c)	
Results :	May explode when heated.
Reliability :	(2) Reliable with restrictions
References :	Budavari, S.ed. The Merck Index - An Encyclopedia of Chemicals,

OECD SIDS	BENZOYL PEROXIDE
2. PHYSICO-CHEN	AICAL DATA ID 94-36-0
	DATE: AUGUST 2002
	Drugs, and Biologicals on CD-ROM version 12:2, Whitehouse Station, NJ, Merck and Co., Inc., 10th edition, 1998
(d) Results Reliability References	 Explodes when heated oxidizer (2) Reliable with restrictions Aldrich, Catalog handbook of fine chemicals, Milwaukee, WI, Aldrich chem. Co., 64, 2000 ~ 2001
(e) Results	: When heated above melting point (103 - 105), instantaneous and explosive decomposition occurs without flame, but the decomposition products are flammable. If under confinement, decomposition may be violently explosive
Reliability References	 (2) Reliable with restrictions Bretherick, L., Handbook of Reactive Chemical Hazards, 4th ed., Butterworths, London, 868 ~ 869, 1990
(f)	
Results	: May decompose explosively if subjected to excessive heat, friction or sudden shock.
Reliability References	 (2) Reliable with restrictions International Agency for Research on Cancer, World Health Organization IARC Monographs on the Evaluation of Carcinogenic Risk of Chemicals to Man, International Agency for Research on Cancer. Geneva, Vol. 36, 267 ~ 283, Supplement 7, p.58
(g)	
Results	 Liable to explode when heated above melting point or when subjected to friction or shock when dry. (2) Reliable with restrictions
References	 (2) Reliable with restrictions Bretherick, L., ed., Hazards in the Chemical Laboratory, 3rd ed., The Royal Society of Chemistry, London, 270 ~ 271, 1981
(h)	
Results	: At elevated temperatures became unstable spontaneously explosive and extremely explosion-sensitive to shock (impact, blows), heat and friction
Reliability References	 (2) Reliable with restrictions National Fire Protection Association, Fire Protection Guide on Hazardous Materials, 7th ed., National Fire Protection Association, Quincy, MA., 49 ~ 113, 49 ~ 114, 491M~57, 491M~145, 1981
(i)	
Results	: Benzoyl peroxide decomposes when heated above 75 (167 F) and will explode when subjected to shock and friction. At elevated temperatures it can be unstable and is spontaneously explosive.
Reliability References	: (2) Reliable with restrictions: U.S. Department of Health and Human Services and U.S.

Department of Labor. NIOSH/OSHA Occupational Health Guidelines for Chemical Hazards, 3 Vol., DHHS (NIOSH) Publication No. 81 ~ 123, January, Volume 1, 1981

2.11 OXIDIZING PROPERTIES

(a) Results Reliability References	 Active oxygen, approximately 6.5 % (2) Reliable with restrictions Lewis, R.J. Sr., ed., Hawley's Condensed Chemical Dictionary, 14th ed., John Wiley & Sons, Inc., New York, 127 ~ 128, 2001
(b) Results	: A powerful oxidizer
References	 (2) Reliable with restrictions Richard, J. Lewis, Sr. SAX' Dangerous Properties of Industrial Materials on CD-ROM 10th ed., John Wiley & Sons, Inc., New York, 2000
(c) Results	: Oxidizer irritant, purity = 75%
Reliability References	 (2) Reliable with restrictions Aldrich, Catalog handbook of fine chemicals, Milwaukee, WI, Aldrich chem. Co., 164, 2000 ~ 2001
(d)	
Results	: An oxidizing agent and strong supporter of combustion
Reliability References	 (2) Reliable with restrictions National Fire Protection Association. Fire Protection Guide on Hazardous Materials. 7th ed. National Fire Protection Association Quincy, MA. 49~113, 49~114, 491M~57, 491M~145, 1981

2.12 OXIDATION: REDUCTION POTENTIAL

No data

2.13 ADDITIONAL DATA

A. Partition co-efficient between soil/sediment and water (Kd)

(a)	
Value	: $K_{oc} = 1,800$
Method	: Determined from an experimental logKow of 3.46 and a
	recommended regression-derived equation
Reliability	: (2) Reliable with restrictions
References	: 1. Online Toxicology Data Network (TOXNET), Hazardous
	Substances Data Bank (HSDB), 2002

2. Lyman, W.J. *et al*, Handbook of Chemical Property Estimation Methods, Washington, DC, Amer. Chem. Soc., $4 \sim 9$, 1990

B. Henry's Law Constant

(a) Value Method Reliability References	 : 0.36 Pa m³/mole : Estimated by HENRYWIN model : (2) Reliable with restrictions : National Institute of Environmental Research (NIER), Korea, Estimation of Physical Chemical Properties and Environmental Fate of SIDS Chemicals (II)-on Benzoyl peroxide, 2002
C. Volatilization from S	oil/Water
(a)	
Result	: Volatilization from moist soil surfaces may be important (1) Volatilization from dry soil surfaces is not expected (2)
Remarks	: (1) Given by estimated Henry's law constant of 3.54×10^{-6} atm m ³ /mole
	(2) Based on estimated vapor pressure of 7.1×10^{-5} mmHg
Reliability	: (2) Reliable with restrictions
References	: 1. Unline Toxicology Data Network (TOXNET), Hazardous Substances Data Park (HSDP) 2002
	 2. Meylan, W.M. and Howard, P.H., Bond contribution method for estimating Henry's law constants, Environ. Toxicol. Chem., 10, 1283 ~ 1293, 1991
(b)	
Result	: Volatilization from water surfaces
Remarks	: Given by estimated Henry's law constant of 3.54×10^{-6} atm m ³ /mole
Reliability	: (2) Reliable with restrictions
Kererences	: I. Unline Toxicology Data Network (TOXNET) : Hazardous Substances Data Bank (HSDB) 2002
	 Meylan, W.M. and Howard, P.H., Bond contribution method for estimating Henry's law constants, Environ. Toxicol. Chem. 10, 1283 ~ 1293, 1991

3.1 STABILITY

3.1.1 PHOTODEGRADATION

(a) Preferred Result		
Туре	: air	
Indirect Photolysis		
Type of sensitizer	: OH radical	
Concentration of sensitizer	: based on intensity of sunlight	
Rate constant (radical)	: $3.55 \times 10^{-12} \text{ cm}^3/\text{molecule-sec}$	
Degradation	ca. 50 % after 3 days (36.1 hours)	
Method	: Estimated by APOWIN model	
Reliability	: (2) Reliable with restrictions	
References	: National Institute of Environmental Research (NIER), Korea,	
	Estimation of Physical Chemical Properties and Environmental	
	Fate of SIDS Chemicals (II)-on Benzoyl peroxide, 2002	
(b)		
Light spectrum	: < 300 nm	
Remarks	: Can decompose on photolysis	
Reliability	: (2) Reliable with restrictions	
References	: International Agency for Research on Cancer, World Health Organization. IARC Monographs on the Evaluation of Carcinogenic Risk of Chemicals to Man, International Agency for Research on Cancer. Geneva, Vol. 36, 267 ~ 283, Supplement 7, p. 58, 1999	
(c)		
Light spectrum	: > 290 nm	
Remarks	: May directly photolyze	
Reliability	: (2) Reliable with restrictions	
References	: Online Toxicology Data Network (TOXNET), Hazardous Substances Data Bank (HSDB), 2002	

3.1.2 STABILITY IN WATER

(a) Preferred Result	
Туре	: Abiotic (hydrolysis)
Half-life at 25 °C	: $t_{1/2}$ at pH 4 = 11.9 hours (Rate Constant, $k_{obs} = 1.62 \text{ l/s} \times 10^5$) $t_{1/2}$ at pH 7 = 5.2 hours (Rate Constant, $k_{obs} = 3.72 \text{ l/s} \times 10^5$) $t_{1/2}$ at pH 9 = Not dectected (Rate Constant, $k_{v} = \text{Not dectected}$)
Degradation %	: The preliminary test at 50 °C for 5 days = 93.5 % at pH 4.0 94.1 % at pH 7.0
	94.2 % at pH 9.0
Method	: OECD TG 111, "Hydrolysis as a Function of pH"
Year	: 2001
10	

OECD SIDS		BENZOYL PEROXIDE
3. ENVIRONMENTAL FA	FE AND PATHWAYS	ID 94-36-0
		DATE: AUGUST 2002
GLP	: No	
Test substance	: Benzoyl peroxide, Source – sigma (Lot No. %	o. 21K0966), purity = 70
	Remarks : Concentration of test substate preliminary test was performed at 50 °C f and 9.0. To test for first – order behave analysed at 25 °C at pH 4.0, 7.0 and 9.0 was determined by HPLC.	ance was 4 mg/l. The for 5 days at pH 4.0, 7.0 for, test substance was Analysis of these tests
Reliabilities	: (2) Reliable with restrictions	
References	: National Institute of Environmental Resea Test of Benzoyl peroxide Hydrolysis as a F LGCI), 2001	rch (NIER), Korea, The Function of pH (tested by

3.1.3 STABILITY IN SOIL

No data

3.2 MONITORING DATA (ENVIRONMENT)

No data

3.3 TRANSPORT AND DISTRIBUTION BETWEEN ENVIRONMENTAL COMPARTMENTS INCLUDING ESTIMATED ENVIRONMENTAL CONCENTRATIONS AND DISTRIBUTION PATHWAYS

3.3.1 TRANSPORT

(a)	
Туре	: Volatility
Media	: Water-Soil
Method	: Estimated volatilization of half-lives
Results	: A model river : 17 days, A model lake : 123 days
Remarks	: (2) Reliable with restrictions
References	: 1. Online Toxicology Data Network (TOXNET), Hazardous
	Substances Data Bank (HSDB), 2002
	2. Lyman, W.J. et al, Handbook of Chemical Property Estimation
	Methods, Washington DC: Amer. Chem. Soc., $15 \sim 1$ to $15 \sim 29$,
	1990

3.3.2 THEORETICAL DISTRIBUTION (FUGACITY CALCULATION)

(a) **Preferred Result**

Media	
Method	
Results	

- : Air-biota-sediment-soil
- : Fugacity level III
- : Benzoyl peroxide would be distributed into environmental compartment; Air (0.04 %), Water (0.02 %), Soil (99.9 %) and Sediment (0.002 %).

Level III				
Compartm ent	Release 100 % to air	Release 100 % to water	Release 100 % to soil	All three
Air	1.08	0.068	0.007	0.04
Water	0.01	85.0	0.0068	0.02
Soil	98.9	6.24	99.999	99.9
Sediment	0.001	8.65	0.0007	0.002

Input parameters

Property	Value			
Chemical Type	1			
Molecular Weight		242.23		
(g/mol)				
Temperature (°C)		25		
Water Solubility		9.1		
(g/m ³)		2.1		
Vapor Pressure (Pa)	0.00929			
Log P _{ow}	3.43			
Melting Point (°C)		105		
	Air	36 (estimated by AOPWIN)		
	Water	5 (derived from hydrolysis in		
Half-life (hour(s))	water)			
	Soil	1E + 11 (default value)		
	Sedime	$1E \pm 11$ (default value)		
	nt	1E + 11 (delault value)		

- Remarks Reliabilities References
- : The EQC model was used for calculation.
- : (2) Reliable with restrictions
- : National Institute of Environmental Research (NIER), Korea, Estimation of Physical Chemical Properties and Environmental Fate of SIDS Chemicals (II)-on Benzoyl peroxide, 2002

(b)

Media Method

- : Air-biota-sediment-soil
- : Fugacity level I

OECD SIDS	BENZOYL PEROXIDE
3. ENVIRONMENTA	AL FATE AND PATHWAYS ID 94-36-0
	DATE: AUGUST 2002
Results	: In air 1.43 %
	In water 28.7 %
	In soil 68.3 %
	In sediment 1.52 %
Remarks	: The EQC model was used for calculation.
Reliabilities	: (2) Reliable with restrictions
References	: National Institute of Environmental Research (NIER), Korea,
	Estimation of Physical Chemical Properties and Environmental Fate
	of SIDS Chemicals (II)-on Benzoyl peroxide, 2002

3.4 IDENTIFICATION OF MAIN MODE OF DEGRADABILITY IN ACTUAL USE

No data

3.5 **BIODEGRADATION**

(a) Preferred Result	
Туре	: Aerobic
Inoculum	: Activated sludge, 30 mg/L as suspended solid
Concentration	: 100 mg/L related to test substance without pre-acclimation
Contact time	: 21 days
Medium	: Water
Degradation	: 83 % after 21 days (based on BOD)
Results	: Readily biodegradable
Kinetic	: 47 % after 7 days (based on BOD)
	83 % after 14 days (based on BOD)
	83 % after 21 days (based on BOD)
	88 % after 21 days (based on TOC)
	100 % after 21 days (based on HPLC analysis)
Method	: OECD TG 301 C, "Ready Biodegradability ; Modified MITI Test
	(I)"
Year	: 1981
GLP	: No
Test substance	: Benzoyl peroxide
Remarks	: Temperature of incubation : $25 \pm 1 ^{\circ}\text{C}$
	300 ml of test solution was made up of 100 mg/L test substance in mineral medium with 30 mg/L activated sludge. Activity control (aniline in mineral medium at 100 mg/L with 30 mg/l activated sludge) and inoculum blank (mineral medium with 30 mg/L activated sludge) were used. Automatic electrolytic biochemical oxygen demand meter for BOD, total organic carbon analyzer for TOC and HPLC were used. BOD was analyzed continuously, TOC and HPLC were measured once after inoculation.
Reliabilities References	 (2) Reliable with restrictions Ministry of International Trade and Industry, Japan, Biodegradation and Bioaccumulation Data of Existing Chemicals Based on the CSCL, 1992

OECD SIDS	BENZOYL PEROXIDE
3. ENVIRONMENTA	L FATE AND PATHWAYS ID 94-36-0
	DATE: AUGUST 2002
(b)	
Туре	: Aerobic
Inoculum	: Adapted ; Secondary activated sludge, domestic
Concentration	: mg/L related to DOC
Contact time	: 28 days
Medium	: Water
Degradation	: 56 % after 28 days (based on the ratio of BOD to ThOD)
Results	: Readily biodegradable
Method	: OECD TG 301D and EEC Part C.6 Degradation-biotic degradation :
	Closed bottle Test
Year	: 1990
GLP	: Yes
Test substance	: Lucidol (Benzoyl peroxide), Source - Akzo Chemicals BV (Lot No.
	589076897), purity = 74.4 %
Remarks	: Since test substance is poorly soluble in water, it was first dissolved
	in dichloromethane.
Reliabilities	: (2) Reliable with restrictions
References	: Ginkel, C.G. van., Akzo Chemicals BV, The Netherlands,
	Biodegradability of Dibenzoyl peroxide, 1990

3.6 BOD5, COD OR RATIO BOD5/COD

No data

3.7 BIOACCUMULATION

(a) Preferred Result	
BCF	: 92
Method	: Estimated by BCFWIN model
Reliability	: (2) Reliable with restrictions
References	: National Institute of Environmental Research (NIER), Korea, Estimation of Physical Chemical Properties and Environmental Fate of SIDS Chemicals (II) -on Benzoyl peroxide, 2002
(b)	
BCF	: 250
Method	: Calculated from experimental logP _{ow} of 3.46 and a recommended regression-derived equation
Reliability	: (2) Reliable with restrictions
References	 1. Online Toxicology Data Network (TOXNET) : Hazardous Substances Data Bank (HSDB), 2002 2. Lyman, W.J. <i>et al.</i>, Handbook of Chemical Property Estimation Methods, Washington DC, Amer. Chem. Soc., 4 ~ 5, 5 ~ 10, 1990

3. ENVIRONMENTAL FATE AND PATHWAYS

3.8 ADDITIONAL REMARKS

Remarks	: From the chemical structure, benzoyl peroxide is expected that it is hydrolyzed to benzoic acid. Benzoic acid is mainly distributed to soil (64.2 %) and water (34.8 %) by Fugacity Level III. No hydrolysis is expected at pH range 4 ~ 11. The substance is readily biodegradable (> 90 % after 28 days) and has a low potential for bioaccumulation
References	: The OECD HPV program, SIDS Initial Assessment Report for 13th SIAM - Benzoates category

4.1 ACUTE/PROLONGED TOXICITY TO FISH

(a) Preferred Result	
Туре	: Flow-through
Species	: Oryzias latipes
Exposure period	: 96 hours
Unit	: mg/L
Analytical monitoring	: HPLC monitoring was performed at 0, 48 and 96 hours.
LC_{50}	: 0.24
Method	: OECD TG 203, "Fish, Acute Toxicity Test"
Year	: 2002
GLP	: Yes
Test substance	: PEROX-B95 (Benzoyl peroxide), Source – Hansol chemience Co. Ltd. (Lot No. 020502), purity = 97.3 %
Remarks	: Test fish ; Age = 9 months / Length = 3.5 ± 0.1 cm / Weight = 0.34 ± 0.04 g
	Test conditions ; Exposure vessel type = 8.7 liter glass aquarium/ Light = $284 \sim 309$ Lux/ Light periocity = $16/8$ hr (light/dark)/DO = $8.0 \sim 8.6$ mg/L/pH = $7.27 \sim 7.55$ /Temperature range = $24.4 \sim 25.0$
	Fish were fed brine shrimp in the morning and tetramin flake in the afternoon.
	Fish were acclimated for 7 days before testing. Dilution water source was tap water which was passed through activated carbon and membrane filter (1 um)
	The water had a hardness of 53.5 mg/L as $CaCO_3$ and an alkalinity of 30.5 mg/l as $CaCO_3$. Test substance was made up to stock solution of 343, 685, 1370, 5193 and 10277 mg/L. 0.15 ml/min of
	stock solution and 200 ml/min of dilution water were supplied to mixing chamber in continuous flow-through test system. Flow rate was 167 ml/min. Dilution water control and acetone control (750 mg/L) were used 10 fish per vessel were exposed to each of five
	test concentrations. Moving-Average Angle was used for statistical method to determine 95 % confidence limits. Nominal concentrations = 0.25, 0.5, 1.0, 2.0 and 4.0 mg/L Measured concentrations = 0.23, 0.47, 0.69, 1.54 and 2.17 mg/L (mean measured concentration)

			5				
		Nominal	Mean measured	Cumulative mortality			
	concentrations		Concentration	(percent mortality)			
		(mg/L)	(mg/L)	24 hr	48 hr	72 hr	96 hr
		Control	Control	0(0)	0(0)	0(0)	0(0)
		Solvent control	Solvent control	0(0)	0(0)	0(0)	0(0)
		0.25	0.23	0(0)	0(0)	0(0)	0(0)
		0.5	0.47	2(20)	6(60)	10(100)	10(100)
		1.0	0.69	4(40)	10(100)	10(100)	10(100)
		2.0	1.54	10(100)	10(100)	10(100)	10(100)
		4.0	2.17	10(100)	10(100)	10(100)	10(100)
Reliabilities References	::	Lowest test sub 0.47 mg/L and was observed in mg/L. Precipita surface were of (1) Reliable wi National Institu Acute Toxicity EG01077, teste	ostance concentrat mortality of contr n 24 and 48 hours ation of test substa oserved at 0.47 - 2 thout restrictions ate of Environmer of Benzoyl perox ed by KRICT), 20	tion caus rols was p at 0.47 p ance and 2.17 mg/l ntal Rese kide to Fi 02	ing 100 % none. Los ng/L and test subs L. arch (NII sh (Repo	% mortali ss of equ l in 24 hr tance at t ER), Koro ort No.	ity was ilibrium at 0.69 he ea, The
(b)							
Туре	:	Semistatic					
Species	:	Poecilia reticul	ata (guppy, local	aquariun	1)		
Exposure period	:	96 hours					
Unit	:	mg/L					
LC_{50}	:	2.0					
Analytical monitoring	:	NO					
Method	:	Vec Vec	and EEC method,	, part C			
GLP Tost substance	:	res Lucidal (Panza	ul porovido). Sou	roo Ala	o Chomi	\mathbf{D}	I at Na
Test substance	·	5589076897), p	urity = 74.4 %	ICE- AKZ	o Chenno	cais DV (LUI NO.
Remarks	:	The range of te mg/L and 10 fis diluting water v ml/L) was use control were al the test medium	st concentration (h (per test concen vas DSW (Dutch d as dispersant. so included. A sh a at all test concen	nominal) tration ar Standard Acetone light pred trations	was 0.7, nd control Water) control cipitate c was obse	, 1.3, 2.4 l) were us and acet and fis on the su rved.	and 4.2 sed. The one (0.1 h water rface of
Reliabilities	:	(2) Reliable wit	h restrictions				
References	:	Mark, U., Meu Acute Toxicity	wsen, I.J.B., Akzo of Dibenzoyl pero	o Chemic xide to fi	cals BV, sh, 1989	The Net	herlands
(c)							
Туре	:	Static					
Species	:	Oryzias latipes					
Exposure period	:	96 hours					
Unit	:	mg/L					

Table showing cumulative mortality :

OECD SIDS	BENZOYL PEROXIDE
4. ECOTOXICITY	ID 94-36-0
	DATE: AUGUST 2002
Analytical monitoring	: No data
LC_{50}	: 3.9
Method	: OECD TG 203, "Fish, Acute Toxicity Test"
Year	: 1996
GLP	: No
Test substance	: No data
Remarks	: Dimethylsulfoxide (DMSO) was used as a dispersant.
Reliabilities	: (2) Reliable with restrictions
References	: Ministry of Environment (MOE), Korea, Establishment of Advanced Testing Methods for Hazardous Chemicals, tested by KRICT, 1996

4.2 ACUTE TOXICITY TO AQUATIC INVERTEBRATES

A. Daphnia

(a) Preferred Result	
Туре	: Static
Species	: Daphnia Magna
Exposure period	: 48 hours
Unit	: mg/L
EC_{50}	: 0.07
Analytical monitoring	: Yes, analytical monitoring was performed by HPLC at 0, 1, 3 and 5
	hours.
Method	: OECD TG 202, part 1, "Daphnia sp., Acute Immobilisation Test"
Year	: 2001
GLP	: Yes

OECD SIDS	ECD SIDS BENZOYL PERC		
4. ECOTOXICITY			ID 94-36-0
		D	ATE: AUGUST 2002
Test substance	: Benzoyl peroxide, So 13174), purity = 79.4	urce-Lancaster Synthesis	Ltd (Lot No.
Remarks	: Test organisms were a Chemistry, Germany group and solvent cor solutions preparations acetone then diluted v	supplied from GSF Institu and juveniles within 24 ho ntrol (Acetone 1000 mg/l) s : 14 mg test substance wa with diluting water.	te of Ecological ours old. Control were used. Stock as dissolved in 10ml
	Test conditions ; Test Exposure vessel type crystallizing dish / Di 211 Annex 2)/Dilutio CaCO3, alkalinity = 3 1427 - 1457 Lux/Ligh chemistry in test and - 8	temperature range = 21.0 = 100 ml test solution in a lution water source = Eler n water chemistry; hardne 39.0 mg/l as CaCO3 ; pH nt periocity = $16/8$ hr (ligh in the control ; DO = 7.5 -	- 21.1 °C / 150 ml hdt M4 (OECD TG ss = 226.5 mg/l as = 8.0/ Lighting = t/dark)/Water 8.6 mg/l, pH = 7.7
	Analytical monitoring hours to identify the the test, contents of l hr. Due to rapid concentrations were system.30 daphnids (exposed to concentration	g was performed by HPI stability of benzoyl perox benzoyl peroxide decrease hydrolysis of benzoyl used for calculating EC 3 replicates : 10 organisme tion of 0.03, 0.06, 0.13, ons)	C at 0, 1, 3 and 5 cide. In the result of ed below 80% for 1 peroxide, nominal C50 value in static s per replicate) were 0.25 and 0.50 mg/l
	Moving-Average Ang determine 95 % confi Cumulative immobili	gle was used for statistical dence limits. zation :	method to
	Nominal	Cumulative number of organ (percent immobility)	iisms immobilized
	Concentration (mg/L)	24hr	48hr
	Control	0(0)	0(0)
	Solvent control	0(0)	0(0)

: 30 Daphnids were immobilized in 0.13, 0.25, 0.50 mg/L at 24 hours and 3 Daphnids in 0.06 mg/L at 48 hours. Added acetone concentration was exceeded 100 mg/L but daphnia response in solvent control was normal.

0(0)

3(10)

30(100)

30(100)

30(100)

0(0)

0(0)

30(100)

30(100)

30(100)

0.03

0.06

0.13

0.25

0.50

OECD SIDS	BENZOYL PEROXIDE
4. ECOTOXICITY	ID 94-36-0
	DATE: AUGUST 2002
Reliabilities	• (1) Reliable without restrictions
References	 (1) Rendole without restrictions National Institute of Environmental Research (NIER), Korea, The acute toxicity of Benzoyl peroxide to Aquatic Invertebrate (<i>Daphnia</i>) (Report No. EG01006, tested by KRICT), 2001
(b)	
Туре	: Static
Species	: Daphnia Magna
Exposure period	: 48 hours
Unit	: mg/L
EC ₅₀	: 2.91
Analytical monitoring	: Yes, measured by NOPC (non - purgeable organic carbon) at 0 hr and 48 hrs.
Method	: OECD TG 202 and EEC Directives
Year	: 1999
GLP	: Yes
Test substance	: Lucidol (Benzoyl peroxide), Source - Akzo Chemicals BV, Purity = 74.4 %
Remarks	: 20 Daphnids divided into 4 batches of 5 animals were used per test concentration and control. The test was carried out as a WAF (Water Accommodated Fraction) with several dilutions. The DSW (Dutch Standard Water) was used as the test medium. The test concentrations were determined by NPOC analysis that was "indicative" or "representative" of test concentrations but not actually measured. These were performed on the undiluted fraction and control.
Reliabilities	: (2) Reliable with restrictions
References	: Wijk, R.J.van and Groenenboom Ir.C.J., Akzo Nobel, The Netherlands, Acute Toxicity of Lucidol to <i>Daphnia magna</i> , 1999

4.3 TOXICITY TO AQUATIC PLANTS

Algae

(a) Preferred Result	
Species	: Selenastrum capricornutum (Strain No. ATCC 22662)
Endpoint	: Biomass, Growth rate
Exposure period	: 72 hours
Unit	: mg/L
E_bC_{50}	: 0.07
E_rC_{50}	: 0.44
Analytical monitoring	: Yes, HPLC monitoring was done at 0 and 24 hours because test substance was not detected after 24 hours.
Method	: OECD TG 201, "Algae, Growth Inhibition Test"
Year	: 2001
GLP	: Yes

OECD SIDS	BENZOYL PEROXIDE
4. ECOTOXICITY	ID 94-36-0
	DATE: AUGUST 2002
Test substance	: Benzoyl peroxide, Source - Lancaster Synthesis Ltd., (Lot No 13174), purity = 79.4 %
Remarks	 No.13174), purity = 79.4 % Laboratory culture of algae was OECD medium and was shook at 100 rpm. Test Conditions ; Test temperature range = 22 - 24 °C/Dilution water source : acetone/Exposure vessel type : 100 ml medium in an Erlenmeyer flask/Water chemistry in test (pH) = pH 7.45 - 7.78 at start (0 hr) and pH 7.45 - 8.01 at end of the test (72 hours), hardness = 226.5 mg/L as CaCO₃, alkalinity = 39.0 mg/L as CaCO₃/Stock solutions preparation : 17 mg test substance was dissolved in 25 ml acetone and diluted with acetone. Light levels and quality during exposure : 8772 - 9106 Lux, Continuously control and solvent control (acetone 1,700 mg/L) were used. Initial cell number of algae was 1 × 10⁴ cells/ml and three replicate vessels were maintained for each treatment and control group. Comprehensive Toxicity Data Analysis and Database Software (Version 5.0) and Dunnett's test (p < 0.05) were used for determination of EC₅₀, LOEC and NOEC value. Nominal concentrations : 0.05, 0.1, 0.2, 0.4 and 0.8 mg/L Cell density at each flask at each measuring point ;

Nominal		Cell density	$(\times 10^4 \text{ cells/r})$	nl)
Concentration (mg/L)	0 hour	24 hours	48 hours	72 hours
Control	1.4	3.1	32	130
Solvent control	1.2	2.4	25	66
0.05	1.3	1.9	22	59
0.1	1.2	2.1	19	67
0.2	1.2	1.5	6.4	53
0.4	0.99	1.1	2.7	20
0.8	1.0	0.78	0.54	0.04

Percent biomass/growth rate inhibition per concentration

Nominal	Growth rates			
Concentration (mg/L)	Growth rates	Relative growth rates (%)	Relative inhibition (%)	
Control	0.063	-	-	
Solvent control	0.055	87.8	12.2	
0.05	0.053	83.8	16.2	
0.1	0.053	84.4	15.6	
0.2	0.052	82.6	17.4	
0.4	0.041	64.4	35.6	
0.8	- 0.013	0	100	

	Growth curves:	8		
	Nominal	Area under the curve		
	Nominal Concentration (mg/L)	Area under	Relative growth	Relative
	Concentration (Ing/L)	the curve	rates (%)	inhibition (%)
	Control	21,708,000	-	-
	Solvent control	13,684,000	63.0	37.0
	0.05	11,984,000	55.2	44.8
	0.1	12,424,000	57.2	42.8
	0.2	7,484,000	34.5	65.5
	0.4	2,752,000	12.7	87.3
	0.8	- 247,600	0	100
Remark Reliabilities References	 Significant difference between test concent concentration was e solvent control was n (1) Reliable without n National Institute of toxicity of Benzoyl p EG01005, tested by F 	ce in the grow tration group an exceeded 100 m normal. restrictions Environmental peroxide to Aqu KRICT), 2001	wth curve was ad each control. A mg/l but growth Research (NIER atic plants (algae	not observed Added solvent n of algae in 2), Korea, The e) (Report No.
(b)				
Species	: Pseudokirchneriella	subcapitata (CC	CAP NO. 278/4)	
Endpoint	: Biomass, Growth rate	e		
Exposure period	: 72 hours			
Unit A nalytical manitoring	: mg/L	analala ananania a	antrain (
Analytical monitoring	performed.	geable organic c	ardon) analysis v	vas
E_bC_{50}	: 0.44			
E_rC_{50}	: 0.83			
Method	: OECD TG 201			
Year	: 1989			
GLP Test substance	: Yes	avida) Carraa	Alma Chamiaala	DV munitar -
l est substance	: Lucidol (Benzoyl per 74.4 %	'oxide), Source-	Akzo Chemicals	BV, purity =
Remarks	: Cultured algae were Fraction) and potass The test concentration "indicative" or "rep actually measured. T and control not done	added to the W ium dichromate ns were determin presentative" of hese were perfo on the diluted fra	WAF (Water - A was used as poned by NPOC and test concentration test concentration the unclassical concentration and the unclassical concentration actions.	accommodated ositive control. alysis that was tions but not liluted fraction
Reliabilities	: (2) Reliable with rest	rictions		
References	: Akzo Nobel, The Net Fraction of Lucidol <i>Pseudokirchneriella s</i>	herlands, Effect on the Growth <i>subcapitata</i> , 198	s of the Water - A of the Freshwate 9	er Green Alga

4.4 TOXICITY TO MICROORGANISMS

(a)	
Туре	: Aquatic
Species	: Secondary activated sludge of a predominantly domestic wastewater
Exposure period	: 30 minutes
Unit	: mg/L
Analytical monitoring	: Yes, the respiration rates of activated sludge were measured in a Biological Oxygen Monitor (BOM)
EC_{50}	: 35
Method	: OECD TG 209 and EEC Part C.Methods for determination of ecotoxicity, biodegradation, activated sludge respiration inhibition test
Year	: 1990
GLP	: Yes
Test substance	: Lucidol (Benzoyl peroxide), Source- Akzo Chemicals BV (Lot No.5589076897), purity = 74.4 %
Remarks	: Dichloromethane (DCM) was used as a dispersant.
Reliabilities	: (1) Reliable without restrictions
References	: Ginkel, van C.G. and Stroo, C.A., Akzo Chemicals BV, The
	Netherlands, Toxicity of dibenzoyl peroxide for activated sludge, 1990

4.5 CHRONIC TOXICITY TO AQUATIC ORGANISMS

4.5.1 CHRONIC TOXICITY TO FISH

No data

4.5.2 CHRONIC TOXICITY TO AQUATIC INVERTEBRATES

No data

4.6 TERRESTRIAL ORGANISMS

4.6.1 TOXICITY TO SOIL DWELLING ORGANISMS

(a)	
Туре	: Artificial soil
Species	: Eisenia foetida
Endpoint	: Mortality
Exposure period	: 14 days
Unit	: mg/kg
LC_{50}	: > 1,000 (dry weight of substance)

OECD SIDS	BENZOYL PEROXIDE
4. ECOTOXICITY	ID 94-36-0
	DATE: AUGUST 2002
Method	: OECD TG 207. "Earthworm. Acute Toxicity Tests"
Year	: 2002
GLP	: No
Test substance	: PEROX-B95 (Benzoyl peroxide), Source – Hansol chemience Co. Ltd (Lot No. 020502), purity = 97.3 %
Remarks	: The limit test of 1,000 mg/kg was shown no mortality.
Reliabilities	: (2) Reliable with restrictions
References	: National Institute of Environmental Research (NIER), Korea, The toxicity of Benzoyl peroxide to earthworm (tested by KRICT), 2002

4.6.2 TOXICITY TO TERRESTRIAL PLANTS

No data

4.6.3 TOXICITY TO OTHER NON MAMMALIAN TERRESTRIAL SPECIES (INCLUDING AVIAN)

(a)	
Species	: Chicken Embryo
Exposure period	: 14 days
Unit	: µmoles/egg
EC ₅₀	: 0.24
$LD_{50}(2d)$: 0.99
LD_{50}	: 0.82
Method	: Other : Embryotoxic effect
Year	: 1984
GLP	: No data
Test substance	: Dibenzoyl peroxide, moistened, purum, Fluka
Remarks	: Embryotoxic effect of benzoyl peroxide to three-day chicken egg was tested. Acetone was used as solvent for benzoyl peroxide and the solution volume of 5 μ l was injected into shell membrane in the air chamber of the egg during 14 days. The eggs were incubated. Malformations, like defects of the right eye of the embroys, were observed.
Reliabilities	: (2) Reliable with restrictions
References	: Korhonen, A., Hemminki, K. and Vainio, H., Embryotoxic Effects of Eight Organic, Peroxides and Hydrogen Peroxide on Three-Day Chicken Embryos, Environ. Res., 33, 54 ~ 61, 1984

4.7 BIOLOGICAL EFFECTS MONITORING (INCLUDING BIOMAGNIFICATION)

No data

4.8 **BIOTRANSFORMATION AND KINETICS**

No data

4.9 ADDITIONAL REMARKS

Remarks	 From the chemical structure, benzoyl peroxide is expected that it is hydrolyzed to benzoic acid. Toxicity of aquatic organisms for benzoic acid is as followed. : Acute toxicity to fish : LC₅₀ (96 hr) of <i>Lepomis macrochirus</i> is 44.6 mg/L and <i>Salmo gairdneri</i> is 47.3 mg/l in static condition. Acute toxicity to aquatic invertbrates : EC₅₀ (48 hr) of <i>Daphnia magna</i> is 100 mg/L in static condition. Acute toxicity to aquatic plants : EC₅₀ (3 h) of <i>Scenedesmus quadricauda</i> is 75 mg/L by inhibition of photosynthesis
References	: The OECD HPV program, SIDS Initial Assessment Report for 13th SIAM - Benzoates category

5.1 ACUTE TOXICITY

5.1.1 ACUTE ORAL TOXICITY

(a) Preferred Result	
Туре	: LD ₅₀
Species/strain	: mouse/ICR
Sex	: male/female
Number of animals	: 5 animals/sex/dose
Vehicle	: 0.5 % methylcellulose sol.
Value	2,000 mg/kg b.w
Route of administration	: oral (gavage)
Method	: OECD TG 401 (acute oral toxicity)
Year	: 2001
GLP	: yes
Test substance	: Benzoyl peroxide, Source $-$ sigma, purity $= 79.9$ %
Test condition	: - Age : 5 weeks
	- Doses : 0 and 2,000 mg/kg
	- Volume administered or concentration : 10 ml/kg
	- Post dose observation period : 14 days
Result	: Mortality : no animals died following exposure.
	Description, severity, time of onset and duration of clinical signs :
	the only clinical symptom observed in 2,000 mg/kg dose group was
	piloerection. After the first day the symptom was recovered in all
	males and female animals.
	Necropsy findings : no abnormal finding was observed in all dosed animals at
	necropsy.
Conclusion	: No death animals were observed at limit test exposed to 2000
	mg/kg . Rat oral $LD_{50} > 2,000 mg/kg b.w$
Reliability	: (1) Reliable without restriction
References	: National Institute of Environmental Research (NIER), Korea,
	Acute oral toxicity of benzoyl peroxide, (Report No. G01083,
	tested by KRICT), 2001
(b)	
Туре	: LD ₅₀
Species/strain	: rats/albino Spartan weighing 203 - 220 grams
Sex	: Male
Number of animals	: 5 animals
Vehicle	: corn oil
Value	: > 5,000 mg/kg b.w
Method	: Other
Year	: 1973

GLP : no

OECD SIDS	BENZOYL PEROXIDE
5. TOXICITY	ID 94-36-0
	DATE: AUGUST 2002
Test substance	: Benzoyl peroxide ; purity = 78 % ; white granules (remainder water)
Test condition	 Groups of 5 male albino rats were administered a single oral dose of 5000 mg/kg of test material in a constant volume (20 ml/kg). Rats were grouped housed. Food and water available ad libitum, except for overnight period prior to administration, during which food was withheld but water was available. Test material was administered via gavage (syringe). All rats were observed for mortality for a period of 14 days. Body weights were measured initially and at 14 days.
Result	: No mortality occurred at the single dose of 5,000 mg/kg. Rats exhibited normal body weight gain. No toxic symptoms or signs were reported. Study was conducted in accordance with a recognized scientific procedure for analyzing the acute oral toxicity of a test material in experimental animals conforming to a Limit Dose study. Although only males were tested and no gross pathology results were reported, the study meets minimum scientific standards and provides sufficient information to support the conclusion that.
Conclusion	: The acute oral LD ₅₀ in male albino rats is greater than 5,000 mg/kg
Reliability	• (2) Reliable with restriction
Remark	 Study was conducted in accordance with a recognized scientific procedure for analyzing the acute oral toxicity of a test material in experimental animals conforming to a limit dose study. However, only males were tested and no gross pathology results were reported.
References	: Wazeter, F.X., and Edwin Goldenthal, Acute Oral Toxicity Test in Male Albino Rats, International Research and Development Corporation (IRDC) August 31, 1973
(c)	
Туре	: LD_{50}
Species	: Rat
Number of animals	: 2 animals/dose
Value	2 > 950 mg/kg b.w
Method	: other
Year	: 1964
GLP	: No data
Test condition	 Benzoyl peroxide, purity = unknown Groups of two fasted rats each were given oral doses of benzoyl peroxide placed on a small amount of pea soup concentration at
Result	 200, 400 and 950 mg/kg. None of the rats died during the study. One of the rats received 400 mg/kg had some vasodilatation, and one that received 950 mg/kg showed slight muscular weakness. The author concluded that the oral LD₅₀ of benzoyl peroxide in rats is greater than 950 mg/kg.

OECD SIDS	BENZOYL PEROXIDE
5. TOXICITY	ID 94-36-0
	DATE: AUGUST 2002
Reliability References	 : (4) not assignable : Toxicity and health hazard summary, report No. 900713. Rochester, NY, Eastman Kodak Co, lab of ind. Med. Health, Safety&Human Factors Lab., 1964, 2p. cited in 'criteria for recommended standard Occupational Exposure to Benzoyl Perovide NIOSH 1977 published'
	reloxide, MOSII 1977 published
(d)	
Туре	: LD_{50}
Species/strain	: Mouse
Value	: 5,/00 mg/kg
Matha d	Discriminating dose : unknown
	: Not specified
ULP Tost substance	: No dala . Ponzovi norovida nurity = unknown
Poliobility	: Delizoyi peroxide, punty – ulikilowii : (4) not assignable
Kenaonnty	No sufficient experimental details
References	 Gigiena Sanitariya, V/O Mezhdunarodnaya Kniga 113095 Moscow, USSR Vol.1, 1936 cited in SAX'S Dangerous Properties of Industrial Materials, Tenth Edition, 1999, John Wiley & Sons, Inc CD version
(e)	
Туре	: LD ₅₀
Species/strain	: Rat
Value	: 7,710 mg/kg
	Discriminating dose : unknown
Method	: Not specified
GLP	: No data
Test substance	: Benzoyl peroxide, purity = unknown
Reliability	: (4) not assignable Markeld LV. Sherrik Varleday Teninelesisheke Varetroni Letek A ringeday
Kelerences	Institut Pro Vychovu Vedoucicn Pracovniku Chemickeho Prumyclu Praha, Czechoslovakia, 1972, cited in SAX'S Dangerous Properties of Industrial Materials, Tenth Edition, 1999 John Wiley & Sons, Inc CD version

5.1.2 ACUTE INHALATION TOXICITY

(a)	
Туре	: LC ₀
Species/strain	: Rat/Spartan albino
Exposure time	: 4 hours
Number of animals	: 10 males ; housed 5/cage
Dose	: 24.3 mg/l
Value	: 24.3 mg/l
Method	: Other (Federal Hazardous Substances Act, Chapter II, Title 16
	CFR.)
Year	: 1973
GLP	: No data

OECD SIDS	BENZOYL PEROXIDE
5. TOXICITY	ID 94-36-0
	DATE: AUGUST 2002
Test substance	: Benzoyl peroxide ; purity = 78 % ; white granules (remainder water)
Test condition	: A single group of 10 male rats were placed in a sealed 59.1 liter glass chamber and exposed for 4 hours to a dynamic atmosphere of test material. Food and water were available <i>ad libitum</i> pre- and post-exposure. Addition of the test material was controlled by two Wright Dust Feeders. Dried, filtered air was passed through the mechanism and directly into the chamber. Airflow was regulated by a flow-meter. Atmospheric concentration was calculated. During exposure rats were separated into 4 units of 2 or 3 each, in order to prevent "piling up", and to insure adequate exposure to each animal. All animals were observed for 14 days following exposure and then sacrificed.
Result	 No mortality was observed at a single dose of 24.3 mg/L. Observable signs/symptoms during the 4 hour exposure period included : eye squint, dyspnea, salivation, lacrimation, erythema, increased and decreased respiratory rates, and increased and decreased motor activity. Observable signs/symptoms after 24 hours post-exposure : all rats appeared normal. Observable signs/symptoms after 48 hours post-exposure included : soft stools. All animals appeared normal from day 5 until the end of the study. Body waight gain was determined to be pormal throughout study.
Reliability	 (2) Reliable with restriction Study was conducted in accordance with a recognized scientific procedure for analyzing the acute inhalation toxicity of a test material in experimental animals. However, only males were tested and no grass pathology results were reported.
References	 Wazeter, F.X., and Goldenthal, E., Acute Inhalation to Rats; 328 - 005 International Research and Development Corporation (IRDC) August 31, 1973

5.1.3. ACUTE DERMAL TOXICITY

No data available

5.1.4 ACUTE TOXICITY, OTHER ROUTES OF ADMINISTRATION

(a)	
Туре	: LD ₅₀
Species/strain	: Mouse/R, CBA
Route of Administration	: i.p.
Exposure time	: Not applicable
Value	: 20 µmoles/mouse (equivalent to 4.84 mg /mouse or 213 mg/kg
	b.w)
Method	: Other
GLP	: No data

OECD SIDS	BENZOYL PEROXIDE
5. TOXICITY	ID 94-36-0
	DATE: AUGUST 2002
Test substance Remarks	 Benzoyl peroxide (recrystallized) Mice were administered 8 different dose conc. of 5, 10, 15, 20, 25, 30, 40, 50 µmoles/mouse. All animals died within 14 days after injection. (2) Poliable with restriction
References	 (2) Reliable with restriction Horgan, V.J., Philpot, J. ST L., Porter, B.W. and Roodyn, D.B., Toxicity of Autoxidized Squalene and Linoleic Acid, and of Simpler peroxides, in Relation to Toxicity of Radiation, Biochem. J., 67, 551 ~ 558, 1957
(b)	
Type Species/strain Route of Administration Exposure time Value	 LD₅₀ Mouse/'R' hybrid, hairless albino i.p. Not applicable 17.1 ± 2.1 (SE) μmoles/mouse (Equivalent to 168 mg/kg b.w - calculated)
Method GLP Test substance Remarks	 Cother Other No data Benzoyl peroxide, purity = unknown Each mouse was administered 10, 15, 20, 30 μmoles of benzoyl peroxide. All animals died within 8 days after injection. The time of death seemed not very dependent on dose.
Reliability References	 (2) Reliable with restriction Philpot, J. St. L. and Roodyn, D.B., A Comparison Between the Effects in Mice of Injected Organic Peroxides and of Whole-body X-Irradiation, Int. J. Rad. Biol., 4, 372 - 382, 1959
(c) Type Species/strain Route of Administration Exposure time Value Method GLP Test substance Remarks Reliability References	 : LD₅₀ : Mouse : i.p. : Not applicable : 250 mg/kg : Not specified : No data : Benzoyl peroxide, purity = unknown : No experimental information is available : (4) not assignable : Takahashi, A., Problems of Hygiene Maintenance for Food Coming into Contact with Rubber and Plastic Products, Nippon Gomu Kyokaishi 48 (9), 537, 1975 [Translated by English, E. A., Internat'l Polymer Sci. Technol., 3 (1), 93 ~ 105, 1976

5.2 CORROSIVENESS/IRRITATION

5.2.1 SKIN IRRITATION

(a)	
Test type	: in vivo
Species/strain	: Rabbits/Albino New Zealand
Results	: Not irritating
Method	: Other
Year	: 1973
GLP	: No data
Test substance	: Benzoyl peroxide, purity = 78 %; white granules (remainder water)
Test condition	 Sex : male/female weighting 2643 - 2952 grams Number of animals : 3 males and 3 females. Total dose : 500 mg of benzoyl peroxide Vehicle : Not used. Exposure time : 4 hours
	Grading scale : Draize scale 500 mg of test material was applied to the abraded and intact sites on the shaved backs of rabbits. It was held in place for 4 hours with a gauze bandage. After 4 hours, the bandages were removed and the exposed areas washed with lukewarm water. The skin was examined for any injury or irritation from benzoyl peroxide at 4, 24 and 72 hours. :
Result	: The skin on the six rabbits appeared unaffected. Due to the absence of skin irritation observed in this study the calculation of primary irritation score was omitted from this reference.
Conclusion	: The authors concluded that 78 % benzoyl peroxide was neither a primary skin irritant nor a corrosive substance.
Reliability	: (2) Reliable with restriction Study was conducted in accordance with a recognized scientific procedure for analyzing the primary dermal irritation of a test material in experimental animals.
References	: Wazeter, F.X., and Goldenthal, E., Primary Skin Irritation and Corrosive Hazard Test in Albino Rabbits, International Research and Development Corporation (IRDC), August 31, 1973
(b)	
Test type	
Species/strain	: Rabbit/Mixed breeding, b.w : 2 - 4 kg
Method	 Slightly irritating Other, 24 hours-patch test (Ziegler, <i>et al.</i>, Dermatol. Monatsschr., 167, 20 ~ 25, 1981; Kligman & Wooding, J. Invest. Dermatol., 49, 78 ~ 94, 1967)
Year GLP	1985 (paper accepted)No data

OECD SIDS	BENZOYL PEROXIDE							
5. TOXICITY	ID 94-36-0							
	DATE: AUGUST 2002							
Test substance Test condition	 Benzoyl peroxide, purity = unknown Sex : Not specified Number of animals : 10 animals/dose Total dose : 0.1, 1, 5, 10, 15 and 30 % of benzoyl peroxide Vehicle : Yellow soft paraffin Control used : negative-vehicle only, positive - 10 % sodium laur sulfate Exposure time period : 24 hours for determining ID₅₀ and 12 da for IT₅₀ Grading scale : No grading system applied 							
	Determining irritant potential - ID_{50} (<i>irritant dose 50</i>) - concentration which causes an irritant reaction (erythematous reaction) after a single epicutaneous application in 50 % of the animals exposed. - IT_{50} (<i>irritant time 50</i>) - number of days necessary to produce an irritant reaction (erythematous reaction) after daily epicutaneous application in 50 % of the animals.							
Remarks Result	 Different concentrations of benzoyl peroxide in yellow soft paraffin were applied under occlusion for 24 hours to shaved areas on the back of rabbits to determine ID₅₀ and IT₅₀. The exposure was repeated once daily for a maximum of 10 days to determine IT₅₀. It was stopped when erythematous reactions occurred. The lowest benzoyl peroxide irritant concentration was 1 % for one animal. All rabbits reacted with 10 %, 15 % and 30 % 							
concentrations without exception. The ID_{50} was determined % of benzoyl peroxide								
	Animal Negative Benzoyl peroxide Sodium lauryl sulfate							

Animal	Negative	В	enzoyl	peroxid	Sodium lauryl sulfate			
number	control	0.1 %	1 %	5 %	10 %	15 %	30 %	10 %
1	_	-	_	+	+	+	+	+
2	-	-	_	+	+	+	+	+
3	-	-	_	+	+	+	+	+
4	-	-	_	+	+	+	+	+
5	-	-	-	-	+	+	+	+
6	-	-	_	-	+	+	+	+
7	-	-	-	+	+	+	+	+
8	-	-	-	+	+	+	+	+
9	-	-	-	+	+	+	+	+
10	-	-	+	+	+	+	+	+
Total	0	0	1	8	10	10	10	10

: The number of days of exposure necessary to produce an irritant reaction in 50 % of the animals with 0.1 %, 1 % and 5 % BP was 8.35 (7.06 - 10.40)d, 4.34 (3.44 - 5.16)d, and 2.31 (1.75 - 2.80)d respectively.

Benzoyl peroxide		Days of exposure										
concentration	1	2	3	4	5	6	7	8	9	10	11	12
0.1 %	0	0	1	1	2	3	3	5	6	6	7	7
1 %	0	0	4	5	6	7	7	8	9	10	10	10
5 %	1	4	8	9	9	10	10	10	10	10	10	10

Reliability References

- : (2) Reliable with restriction
- : Haustein, U-F, Tegetmeyer, L. and Ziegler, V., Allergic and Irritant Potential of Benzoyl Peroxide, Contact Dermatitis 13, 252 ~ 257, 1985

1	-	λ.
(с)
١.	v	,

Species/strain Results Method	Guinea pigSlightly irritatingOther
GLP	: no
Test substance	: Benzoyl peroxide, purity = unknown
Kemarks	: Patches of skin were chemically depliated, and pure benzoyl peroxide, in doses ranging from 0.25 to 1.0 g/kg, was held against the depilated skin under patches for 24 hours. The skin under the benzoyl peroxide was examined for any irritation or other injury. Slight erythema with some delayed scarring of the epidermis resulted. There were no deaths. A similar test was run on guinea pigs with 10 % solution of benzoyl peroxide in propylene glycol. The doses ranged from 5 to 20 ml/kg. Only slight erythema was observed ; no deaths occurred.
Reliability	: (4) not assignable Experimental details are insufficient.
References	: Toxicity and Health Hazard Summary, report No. 900713, Rochester, NY, Eastman Kodak Co, laboratory of Industrial Medicine, Health, Safety, and Human Factors laboratory, 1964, 2 pp cited in 'criteria for recommended standard Occupational Exposure to Benzoyl Peroxide, NIOSH 1977 published'
(d) Species/strain Results Method GLP	 Rabbit/New Zealand white Irritating Other (cumulative irritation study) No data

OECD SIDS			BENZOYL PEROXIDE						
5. TOXICITY	ID 94-36-0								
			DATE: AUGUST 2002						
Test substance	: Tritium benz radiochemic	: Tritium benzoyl peroxide, purity = 97 %; chemical and radiochemical purity							
Remarks	: The study c topical form dispersed o emulsion v formulations backs of rat daily with th 0.5, barely p even eryther intense eryt erythema pl	 radiochemical purity The study compared the skin irritation of benzoyl peroxide from topical formulations in which the substance was either freely dispersed or was entrapped in a SOWE (simple oil-in-water emulsion vehicle). Two groups of rabbits received different formulations. 0.025 ml of each formulation was applied to clipped backs of rabbits for 21 days. Irritancy observations were recorded daily with the following scale : Grade 0, no skin reaction ; grade 0.5, barely perceptible, spotty erythema ; grade 1.0, mild, even erythema covering the whole area of application ; grade 2.0, intense erythema with clearly defined edge ; grade 3.0, intense erythema plus edema, raised edge ; grade 4.0. intense ervthema. 							
	edema, and formulation 21 days. Cur	l bullae ; grade 5.0, nec was applied over a previou mulative 21- day irritation so	rosis. 0.025 ml of each sly marked 2 cm ² area for cores were as follow ;						
		5	, 						
		Free bezoyl peroxide	Entrapped benzoyl peroxide						
	vehicle	1	3						
	2.5 % BP	280	160						
	5.0 % BP	520	150						
Reliability References	 This study s skin irritatin (2) Reliable Wester, R.C Maibach, H Porous Mid Irritancy, J. A 	 This study shows the entrapped/controlled-released system is less skin irritating. (2) Reliable with restriction Wester, R.C., Patel, R., Nacht, S., Leyden, j., Melendres, J. and Maibach, H., Controlled Release of Benzoyl Peroxide from a Porous Microsphere Polymeric System Can Reduce Topical Irritancy, J. Am. Acad. Dermatol., 24, 720 ~ 726, 1991 							
(e)									
Species/strain	: Mouse/albin	10							
Results	: Slightly irrit	ating							
CLP	: Other (cumi	lative irritation study)							
ULF Test substance	• 10 % Benzo	vl perovide							
Remarks	 Approximat entire dorsur histological marked va degenerating over the enir 	ely 100 μl of benzoyl per m for 6 months. Full thickne microscopic examination. riations in size and sh g and separating each other dermis. Dermal changes wer	roxide was applied to the ess excisions were made for The keratinocytes showed ape (atypia). Some are the A focal serous crust lies e not notable						
Reliability References	 (2) Reliable Kligman, A Assessing the Food and Ch 	with restriction .M. and Kligman, L.H., A he Chronic Toxicity of Top nemical Toxicology, 36, 867	Hairless Mouse Model for bically Applied Chemicals, ~ 878, 1998						

5.2.2 EYE IRRITATION/CORROSION

(a)											
Test type	: <i>in vivo</i> eye irritation study	: <i>in vivo</i> eye irritation study									
Species/strain	: Rabbit/New Zealand White	Rabbit/New Zealand White									
Method	: OECD TG 405 (1987)	OECD TG 405 (1987)									
Year	: 2001	2001									
GLP	: Yes										
Test substance	: Benzoyl peroxide, min., pu	rity = '	70 %,	rema	ainde	r wa	ter				
Test condition	: - Sex : male										
	- Number of animals per do	ose : 4	anima	ls/dc	se						
	- Dosed used : 100 mg/site	/rabbit									
	- Observation period : from	n 1 hou	r aftei	dos	e to 7	/2 hc	ours				
	- Tool used to assess score	: Hand	-slit la	amp							
	- Scoring method used : Ol	- Scoring method used : OECD method									
	- Vehicle and eye washing	- Vehicle and eye washing : eye washed by saline solution at 30									
	seconds after instillation for	r 3 ani	mals.	No v	vashi	ng fé	or 1 a	anima	al		
Result	: Corrosiveness was not obse	: Corrosiveness was not observed. Slight symptoms of irritation were									
	observed when 100 mg of	`benzo	yl pe	roxid	e ins	stille	d to	an ey	ye of		
	rabbit.										
	- Irritation Score (Cornea	- Irritation Score (Cornea/Iris) :									
	Eye response	Cor	mea			Iris					
	Observation time	1	24	48	72	1	24	48	72		
	M1(No eye washed animal)	1	1	1	0	1	1	1	0		
	M2	0	0	0	0	0	0	0	0		
	M3	0	0	0	0	0	0	0	0		

The area of corneal opacity : < 1/4

M4

- Irritation Score of Conjunctivae (Redness/Chemosis) :

0

0

0

0

0

0

0

0

Eye response	Redr	Redness Chemosis						
Observation time		24	48	72	1	24	48	72
M1(No eye washed animal)	2	2	1	1	2	2	1	0
M2	1	0	0	0	0	0	0	0
M3	1	1	1	0	0	0	0	0
M4	0	0	0	0	0	0	0	0

OECD SIDS	BENZOYL PEROXIDE
5. TOXICITY	ID 94-36-0
	DATE: AUGUST 2002
Reliability References	 clinical signs : animal survived during the test and no adverse clinical symptom was observed. No eyes washed animal : opaque cornea was observed until 48 hours after dose in and its area was less than 1/4. Deepened rugae were also found in iris till 48 hours after dose. In the conjunctivae, obvious redness, swelling and excrement were observed till 48 hours after dose but completely recovered after 72 hours. Eyes washed animals : No significant clinical signs were observed. (1) Reliable without restriction National Institute of Environmental Research (NIER), Korea, Acute Eye Irritation/Corrosion Test in Rabbits (Report No. S367, tested by LGCI), 2002
(b)	
Test type	: <i>in vivo</i> eye irritation study
Species/strain	: Rabbit/Albino New Zealand
Method	: Other (Federal Hazardous Substances Act, Chapter II, Title 16 CFR, US)
Year	: 1973
GLP	: No data
Test substance	: Benzoyl peroxide, purity = 78 %; white granules (remainder water)
Test condition	 Sex : male and female Number of animals : 4 animals/sex. 5 rabbits in Group I and 3 rabbits in Group II. Dosed used : 0.1 ml of test material was instilled into the everted lower eye lid of one eye of each rabbit. The average weight of the test material was 111.4 (group I, 5 minute wash) and 120.7 mg (group II, 24 hours wash). Exposure time : group I - 5 minutes, group II - 24 hours. irritation score : 0-4 Observation period : Eyes were examined at 1, 24, 48, 72 hours and 7 days after washing.
Result	 From I : benzoyl peroxide produced very minor redness (conjunctivitis) and swelling (chemosis) in 1/5 and 2/5 rabbits, respectively, but it was clear in 24 hours. No corneal opacity was observed. Group II : there was grade 1 opacity in 3/3 at 24 hours ; clear in 3/3 at 48 hours. No iritis. There was grade 2 redness in 2/3 at 48 hours, degrading to grade 1 in 1/3 at 72 hours : clear by day 7
Conclude	 Benzoyl peroxide was neither irritating nor corrosive to the eyes of albino rabbits if it was washed out within 5 minutes, but if 78 % benzoyl peroxide was no t washed out until 24 hours later, it proved to be a strong irritant. It was not considered corrosive because corneal opacity lasted less than 6 days.
Reliability	: (2) Reliable with restriction

OECD SIDS	BENZOYL PEROXIDE
5. TOXICITY	ID 94-36-0
	DATE: AUGUST 2002
References	: Wazeter, F.X. and Goldenthal, E., Eye irritation in Albino Rabbits, International Research and Development Corporation (IRDC), August 31, 1973
(c)	
Species/strain	: Rabbit
Method	: Other
GLP	: No data
Test substance	: Benzoyl peroxide, purity = 50 and 93 %; powder
Remarks	: The eyes were rinsed with water 1 minute after instillation, and any solid residues were removed with a cotton swab.
References	: Kuchle, H.J, Untersuchungen über die augenschädigende Wirkung organischer Peroxyde, Zentralblatt für Arbeitsmedizin und Arbeitsschuntz, Heit 2, 25 ~ 31, 1958

5.3 SKIN SENSITIZATION

(a) Type Species/strain Method Year GLP Test substance	 Cellular proliferation Mouse, CBA/Ca strain or CBA/JHsd strain Local lymph node assay (draft OECD TG429) 1997 (paper received) No data Benzoyl peroxide (CAS No. 94-36-0) Source - Sigma, USA, purity = 70 %
Test condition	 Sex : Female Age of animal at study initiation : 6 - 12 weeks Number of animals per sex per dose : 5 animals/dose Route of administration : topical application on the dorsum of both ears Induction concentration : 0.5, 1.0, 2.5, 5.0 and 10.0 % benzoyl peroxide Induction vehicle : Acetone Negative control : vehicle only Volume of material dosed : 25 µl Duration of exposure for induction : daily for 3 consecutive days. Length of rest period : 2 days prior to analysis Grading system used : <i>in vivo</i> [³H]methylthymidine(or [¹²⁵I]iododeoxyuridine) : incorporation into lymph node cell DNA associated with proliferation induced by application of benzoyl peroxide was an objective and quantifiable response. Stimulation indices (SI) were determined as increase in ³H - TdR incorporation relative to vehicle-treated controls. SI of 3 or greater was considered to have skin sensitizing activity. Based on fitted quadratic regression analyses, estimates of the applied concentration of benzoyl peroxide required to elicit a SI of 3 (EC 3 value) were calculated.

OECD SIDS 5. TOXICITY

Result

: Even at lowest concentration tested (0.5 %), benzoyl peroxide induced stimulation indices of considerably greater than 3, and for this reason it proved impossible to derive mathematically EC_3 values for this chemical.

	Lab. A	Lab. B	Lab. C	Lab. D	Lab. E
Concentration	dpm (SI)	dpm (SI)	dpm (SI)	dpm (SI)	dpm (SI)
Actione	307	390	463±36	262±65	27±6
0.5	4457	9110	8640±953	3581±851	658±86
	(14.6)	(23.4)	(18.7)	(14.7)	(24.4)
1.0	5273	8880	9746±1192	2061±956	598±81
	(17.2)	(22.8)	(21.0)	(7.9)	(22.1)
2.5	5555	8495	11543±1909	2859±1017	909±129
	(18.1)	(21.8)	(24.9)	(10.9)	(33.7)
5.0	6189	8762	11474±1422	5373±1176	849±56
	(20.2)	(22.5)	(24.8)	(20.5)	(31.4)
10.0	6681	6274	8595±971	4522±409	715±62
	(21.8)	(16.1)	(18.6)	(17.3)	(26.5)

- stimulation index(SI)

All concentrations of benzoyl peroxide elicited a statistically significant increase in isotope incorporation in the 3 laboratories (lab C, D, E)

Conclusion	: Benzoyl peroxide provoked very vigorous sensitising responses at all test concentration.
Reliability	: (1) Reliable without restriction
5	This substance was evaluated in each of five independent international laboratories using either the standard LLNA protocol (Kimber & Basketter, 1992) or minor modifications of it.
References	: Kimber, I., Hilton, J., Dearman, R.J., Gerberick, G.F., Ryan, C.A., Basketter, D.A., Lea, L., House, R.V., Ladics, G.S. Loveless, S.E. and Hastings, K.L., Assessment of the Skin Sensitization Potential of Topical Medicaments Using the Local Lymph Node Assay: An Interlaboratory Evaluation, J. Toxicol. Environ. Hlth., Part A, 53 (7), 563 ~ 579, 1998
(b)	
Туре	: Challenge (Buehler test)
Species/strain	: Guinea pig
Number of test animals	: 20 test animals in the test group, 10 naive control animals for challenge and 10 separate naïve control animals for rechallenge
Route of administration	: topical with occluded patch
Results	: Sensitizing
Method	: Guinea Pig Maximisation Test (OECD TG 406)
GLP	: No data
Test substance	: Benzoyl peroxide, purity = unknown
Result	: 42 % of the test group was judged as having positive allergic reactions. It is classified as skin sensitizer according to the threshold defined by EC (> 15 %).

OECD SIDS	BENZOYL PEROXIDE
5. TOXICITY	ID 94-36-0
	DATE: AUGUST 2002
Reliability	(2) Reliable with restriction
References	Basketter, D.A., Gerberick, G.F., An interlaboratory evaluation of the Buehler test for the identification and classification of skin sensitizers, Contact Dermatitis, 35, 146 ~ 151, 1996
(c)	
Туре	Challenge
Species/strain	human/white young
Method	Other, Maximization test
GLP	No data
Test substance	Benzoyl peroxide (topical preparations for acne)
Test condition :	4 formulation of benzoyl peroxide, two 5 % and two 10 % gel, were evaluated in a group of 50 white college students. The subjects were challenged 10 - 14 days later by a 48 hours patch test and the responses compared to pretesting with the same materials. The challenge site was evaluated at 48, 72, and 96 hours on a four point scale
Pagult	point scale. 28 of 50 subjects developed $a \pm a$ or greater reaction. Sensitized
Result	subjects reacted to all four formulations
Reliability	(2) Reliable with restriction
References	Leyden, J.J., Kligman, A.M, Contact sensitization to benzoyl peroxide, Contact Dermatitis, Vol 3, 273 ~ 275, 1977
(d)	
Туре	TINA test
Species/strain	Guinea pig/albino
Results	Sensitizing
Method	Other
GLP	No data
Test substance	Benzoyl peroxide, purity = unknown
Remarks	5 of 25 guinea pigs had a positive reaction after the TINA procedure on 2 consecutive days, while the other animals and controls were negative. The proportion of 20 % positive reactions to Benzoyl peroxide can be interpreted as indicating a relatively weak sensitization rate.
Reliability	(2) Reliable with restriction
References	Haustein, U-F, Tegetmeyer, L. and Ziegler, V., Allergic and Irritant Potential of Benzoyl Peroxide, Contact Dermatitis, 13, 252 ~ 257, 1985

5.4 REPEATED DOSE TOXICITY

(a) Preferred Result									
Species/strains	:	: Rat/Sprague-Dawley							
Sex	:	: Male/Female							
Route of administration	:	Oral (gav	/age)						
Method	:	OECD	TG	422	(Combined	Repeated	Dose	and	
		Reprodu	ction/D	evelopn	nental Toxicity	Screening Tes	st)		
Year	:	2001		-	-	-			
GLP	:	Yes							

OECD SIDS	BENZOYL PEROXIDE								
5. TOXICITY	ID 94-36-0								
	DATE: AUGUST 2002								
Test substance Dose level Exposure period Frequency of treatment Control group Post exposure observation period Statistical methods	 Source - SIGMA, purity = 70 % 0, 250, 500, 1,000 mg/kg b.w/day Male : 29 days, female : from 2 weeks before mating to the day 3 of lactation Daily (-) Concurrent vehicle (+) Cyclophosphamide (4.5 mg/kg/day) 1 day Dunnett's multiple comparison test 								
Test condition	 : - Test Subjects Age : 7 weeks old for male and female No. of animals : 10 animals/sex/dose (5 animals/sex/positive control) Study Design Vehicle : Corn oil Satellite groups : Recovery group (5 animals/sex) at 0 and 1000 mg/kg for follow-up observation. Clinical observations performed and frequency : General condition and body weight were observed once a day and once a week respectively. For pregnant females, body weight was determined on the day 1, 8, 15 and 20 of gestation and 1 and 3 of lactation. Food consumption was determined on the day 8, 15 and 29 of exposure. Haematology and biochemistry for males only at time of necropsy after 29 days of chemical exposure. Organ weight : liver, kidney, adrenal gland, testis, epididymis, thymus, spleen, brain, heart, ovary, uterus, thyroid gland, spermary, prostrate gland Microscopic : brain, spinal cord, stomach, pancreas, jejunum, ileum, cecum, colon, rectum, liver, kidney, adrenal gland, spleen, heart, thymus, thyroid, bronchus, lung, pituitary gland, ovary, uterus, vagina, testis, epididymis, spermary, prostrate gland, mammary gland, bladder, nodi lymphatici mesenterici, nodi lymphatici, nervus ischiadicus, femoral marrow and all gross lesion. 								
NOAEL LOAEL	 500 mg/kg b.w/day 1,000 mg/kg b.w/day 								
5 TOXICITY						-	BENZO	YL PER	OXIDE
-------------	---	---	---	---	--	--	--	---	---
5. 10/10/11							DATE:		94-36-0 ST 2002
Results	: - Body we (approximate exposed to weight char recovery gre- - Food/wate mg/kg was chemical e showed abn - Description : Two femate to 500 mg diarrhea an Hematemess are not co exposure.	eight tely 7 1,000 nges v oup. er con signifi xposu ormal or, sev le rats /kg. T d urin is was nsider	: Boo % mo mg/kg vere of sumpta focantly re in food c erity, t of dep Wo fe e were s seen ed to	dy we re than g only oserved ion : T increa male. consum ime of pilation emale of e show in one have	ight g contr after l in all he meased sed du No c ption. <i>onset</i> and cr of ger ed in male been	gain w ols) in f 29 days 1 anima an food uring fir other tr and dur tust wer tital con 1,000 k at 500 r related	as slight female in s. No si ls except consum- st and for eatment <i>ration o</i> re found ntamina ag/mg ra- mg/kg.	htly ind recovery gnifican ot highe nption o ourth we groups <i>f clinica</i> when e tion ow ecovery These finizoy	creased group at body st dose f 1,000 eeks of s were al signs xposed ving to group. indings eroxide
	- Hematolo	oical d	and bi	ochem	ical fii	ndinos i	nciden	re and s	everity
	:	Sicur		<i>circini</i>	icut jti			i unu s	everity
	Hematologi concentratio group. Slig recovery gr any other an Biochemica significant observed an - <i>Histopath</i> treatment microscopio change in multinuclea affects were	cal fir on of htly coup. N nimals l find decrea d that ology group cally a sperm ted gi e obser	idings male a lecreas lo othe ings : ase in also sl <i>incid</i> o, tea and th cell ant cell ved in	: sligh and fer se in h er hema in fer aspar howed <i>ence a</i> st is ese pa such a ll. In th uterus	tly inc male r nemato atologi male tate a in 1,00 md se deg thosis is apo ne fem expos	rease in ats in ocrit in ical cha rats rea minotra 00 mg/k <i>verity</i> : eneration were p ptosis, ale repr ed to 1, <u>o mean</u>)	n RBC a 1,000 m 1,000 nges we ceiving unsperas g recove In the ons we mainly cellular coductiv 000 mg/	and hen ng/kg re mg/kg ere obse 250 m ee (AST ery grou e 1,000 ere ob morpho swellin re organ /kg.	hatocrit ecovery female rved in g/kg a T) was up. mg/kg observed ological ng and , slight
	Treatment	TP	ALB	AST	ALT	BUN	CREA	O TCH	GLU
	(mg/kg/day)	g/dL	g/dL	IU/L	IU/L	mg/dL	mg/dL	mg/dL	mg/dL
	0	6.9	2.5	145.5	51.1	18.0	0.6	90.8	87.0
	250	6.7	2.5	88.2	59.6	17.0	0.6	90.4	87.4
	500	6.3	2.3	122.6	56.9	18.9	0.6	101.4	82.8
	1,000	6.4	2.4	102.9	61.3	18.3	0.5	92.4	98.6
	recovery	7.5	2.9	118.8	26.9	20.8	0.6	80.8	112.6
	-		1	1	1 0 7 0		i	-	-

- *Mortality and time to death* : No deaths were found in all animals including control groups.

- Ophthalmologic findings : Not examined

- Organ weight changes

mean organ weight of male : Significant decrease in absolute and relative weights of left testes and relative wt of epididymis were observed in 1,000 mg/kg exposure group (p < 0.05), however these changes were not seen in recovery group. No other significant organ changes were found in all male rats.

Dose (mg/kg)	0	250	500	1,000	(+)	satellite
LT Testes (g)	1.714	1.645	1.686*	1.496	1.582	1.592
RT Testes (g)	1.734	1.638	1.684	1.589	1.585	1.582
LT Epididymis (g)	0.633	0.609	0.595	0.553	0.578	0.65
RT Epididymis (g)	0.648	0.611	0.595	0.576	0.597	0.66
Liver (g)	1.396	11.887	12.193	11.985	11.299	12.112
Thymus (g)	0.441	0.437	0.437	0.453	0.315	0.437
LT Kidney (g)	1.451	1.419	1.646	1.482	1.468	1.51
RT Kidney (g)	1.476	1.448	1.609	1.521	1.46	1.47
LT Adrenal (g)	0.035	0.032	0.035	0.038	0.032	0.036
RT Adrenal (g)	0.032	0.032	0.034	0.038	0.028	0.033
Spleen (g)	0.765	0.742	0.821	0.77	0.526	0.78
Brain (g)	2.09	2.058	2.033	2	2.066	1.917
Heart (g)	1.419	1.352	1.389	1.295	1.401	1.327

mean organ weight of female : increase in absolute and relative liver weights at 1,000 mg/kg, but decrease in relative liver weight at 1,000 mg/kg recovery group.

Dose (mg/kg)	0	250	500	1,000	(+)	satellite
Liver (g)	9.088	10.124	10.013	10.15	7.435	7.086
LT kidney (g)	0.878	0.997	0.907	1.023*	0.846	0.832
RT kidney (g)	0.913	0.989	0.929	1.059*	0.868	0.851
LT adrenal (g)	0.039	0.04	0.039	0.044	0.036	0.03
RT adrenal (g)	0.034	0.039	0.037	0.04	0.035	0.03
Thymus (g)	0.244	0.345	0.377	0.342	0.211	0.342
Brain (g)	1.882	1.96	1.924	1.982	1.912	1.878
Spleen (g)	0.579	0.659	0.604	0.662	0.413	0.558
Heart (g)	1.029	0.965	1.022	1.004	1.068	0.874

- Histopathology incidence and severity : In the 1,000 mg/kg treatment group, testis degenerations were observed microscopically(p < 0.05). In the female reproductive organ, slight effects were observed in uterus exposed to 1000 mg/kg.

Male

Treatment (mg/kg/day)	0	250	500	1,000	(+)
No. of animals	10	10	10	10	5
Testes					
Degeneration					
Minimal				5*	2
Marked				1	
Epididymidis					
Degeneration				2	1
Minimal				5	1
Marked				1	
Decrease of spermatocyte				1	

Treatment (mg/kg/day)	0	250	500	1,000	C (+)
<u>Kidney</u>					
Congestion					
Minimal	1	3			1
Mineralization					
Focal, minimal	1	2	2		2
Focal, slight				2	
<u>Uterus</u>					
Epithelial vacuolation					
Minimal			1	1	
Epithelial hyperplasia					
Minimal			1	1	
Inflammatory cell					
infiltration					
Focal, minimal				1	

Conclusion

Reliability

: Significant decrease in absolute and relative weights of left testes and relative wt of epididymis were observed with testis degenerations in 1,000 mg/kg treated group. This result indicated the NOAEL is 500 mg/kg to male rats.

- : (1) Reliable without restriction
- References : National Institute of Environmental Research (NIER), Korea, Combined Repeated Dose Toxicity with the Reproduction/Developmental Toxicity Screening Testing of Benzoyl peroxide in Rats, (Report No.P049, tested by LGCI), 2001

OECD SIDS	BENZOYL PEROXIDE
5. TOXICITY	ID 94-36-0
	DATE: AUGUST 2002
(b)	
Species/strains	: Rat and mouse/albino
Sex	: Male/Female
Route of administration	: Feeding (p.o) and dermal (subcutaneous injection, skin painting)
Method	: Other (chronic)
Year	: 1964
GLP	: No data
Exposure period	: 80 (mice) and 120 (rat) weeks
Frequency of treatment	: Daily
Dose	: 0.2826, 0.02826 and 0.002826 % in diet
Control group	: No treatment and vehicle only control group
NOEL	: 2.8 ppm
LOEL	: N/A
Test substance	: Novaldelox (commercial powder containing 18 % Benzoyl peroxide)
Test condition	 4 experimental groups (FC, F1, F2 and F3) were maintained on a diet of whole meal flour containing 0, 28.26, 282.6, 2826 ppm of benzoyl peroxide. 25 animals /species/dose/sex were used. Other 3 experimental groups (BC, B1 and B2), composed of 100 and 25 animals/species/dose/sex, were given breadcrumbs made from flour containing 28.26 and 2.826 % benzoyl peroxide. 2 groups (SIC, SI) were given subcutaneous injection of 0, 50, 120 mg benzoyl peroxide. Another one group (T), consisting of 25 male and female mice, were painted on the back of the neck on 6 days each week with one drop (50 mg approximately) of 50 % suspension of benzoyl peroxide in flour paste. Group SIC, SI, T and TC were provided with a commercial pellet diet. Body wt gains of rats were noted weekly for 18 weeks, and then monthly. The animals were examined twice daily with regard to their general health. Animals that were moribund were killed. All animals that were killed or that died were autopsied.
Kesuits	: The rats consuming the flour diets containing the two highest levels of benzoyl peroxide showed significant decrease in body wt. gain. There were no significant differences in the body weight of the rats given treated breadcrumbs. There was no significant difference between experimental and appropriate control groups in the mortality rate of the animals except F2, in which there was a large number of accidental deaths. There were statistically significant incidence of atrophy of the testicles in the rats given the diet based on flour treated with benzoyl peroxide at 2826 ppm and in the rats receiving diets of breadcrumbs made with flour treated with benzoyl peroxide at 28.26 and 2.826 ppm.

Infective	F1	F2	F3	FC	B1	B2	BC
Degenerative	138	109	136	159	110	130	104
Vascular including	53	62	49	48	49	49	56
haemorrhage	16	13	15	28	27	23	49
Miscellaneous	38	30	32	41	41	47	34
Tumor (ben., mal.)	12	2	3	9	31	11	35
Ulceration of stomach	2.2	-	-	-	1.5	2.1	-
Atrophy of testes	27*	15	15	4	19*	23^{*}	11

- Pathological finding in rats

* Significant difference from controls P < 0.05

The authors suggested that this atrophy was caused by benzoyl peroxide, which probably marginally decreased the amount of vitamin E in the diet.

The injection of benzoyl peroxide subcutaneously in rats and mice did not result in any tumor formation at the site of injection, or in any increase in tumor incidence elsewhere. Application of a paste made from flour treated with benzoyl peroxide to the skin of rats and mice also caused no tumor formation.

Reliability	: (2) valid with restriction
Remark	: Exact doses cannot be determined since the actual diet consumption
	was not reported.
References	: Sharratt, M., Frazer, A.C. and Forbes, O.C., Study of the Biological
	Effects of benzoyl Peroxide, Fd. Cosmet. Toxicol., Vol.2, 527 ~ 538,
	1964

5.5 GENETIC TOXICITY IN VITRO

A. BACTERIAL TEST

(a) Preferred Result

Туре	: Bacterial reverse mutation assay
Species/Strain	: Salmonella typhimurium TA 98, TA 100, TA 1535, TA 1537, Escherichia coli WP2 uvrA
Method	: OECD TG 471, 472
System of testing	: Bacterial
Year	: 2001
GLP	: Yes
Metabolic activation	: - Species and cell type : Rat Sprague Dawley strain, male, liver homogenate
	- Quantity : 4 % S9 in the S9 mix
	- Induced : Aroclor-1254 induced
Concentrations tested	: 0.247, 0.74, 2.22, 6.67, 20 and 60 µg/plate
Statistical Methods	: Dunnett's test $(p < 0.01)$
Test substance	: Benzoyl peroxide, source – Sigma (B-2030), Purity = approx. 71.8 %

OECD SIDS	BENZOYL PEROXIDE
5. TOXICITY	ID 94-36-0 DATE: AUGUST 2002
Test condition	 Number of replicates : one time Frequency of Dosing : 3 plates/dose Positive and negative control groups and treatment : Negative control- solvent control (Dimethyl sulfoxide), Positive control - 2-aminoanthracene with S9, 2-nitrofluorene, sodium azide, 9-aminoacridine, 4-nitroquinoline 1-oxide Number of metaphases analyzed : not analyzed Solvent : Dimethyl sulfoxide Description of follow up repeat study : A preliminary test was performed at the concentrations of 1.6, 8, 40, 200, 1,000 and 5,000 µg/plate. Criteria for evaluating results : counting the number of revertant colonies in the plate after 2-day-incubation at 37 celcius.
Cytotoxicity conc	: With metabolic activation : not observed
Genotoxic effects	Without metabolic activation : not observed With metabolic activation : negative
	Without metabolic activation : negative
Conclusion	: Benzoyl peroxide did not induce mutation in the Salmonella typhimurium TA98, TA100, TA1535, TA1537, Escherichia coli WP2 uvrA strains with or without metabolic activation.
Reliability	: (1) Reliable without restriction This study was well conducted by GLP laboratory based on OECD test guideline.
References	: National Institute of Environmental Research (NIER), Korea, Bacterial Reverse Mutation Test, Test No. S350, tested by LGCI, 2001
(b)	
Туре	: Bacterial reverse mutation assay
System of testing Concentration Metabolic activation Results	 Salmonella typhimurium TA 97, TA102 0.001, 0.005, 0.01, 0.05, 0.1 μg/plate With and Without
Cytotoxicity conc	: With metabolic activation : not observed Without metabolic activation : slight cytotoxicity was observed at the doses of 0.1 mg/plate and 0.05 mg/plate. Number of revertant colonies was reduced to 40 % in TA97 and to 24 % in TA102 strain.
Precipitation conc Genotoxic effects	 With metabolic activation : negative Without metabolic activation : negative
Method	: Other (Preincubation method modified by Ames)
GLP	: No data
Test substance	: Source - Wako Pure Chemical Industries, Lot No. DCE 2342, Japan.
Reliability	: (2) Reliable with restriction
Keterences	: Fujita, H. and Sasaki, M., Mutagenicity Test of Food Additives with <i>Salmonella typhimurium</i> TA 97 and TA102 (II), Annu. Rep. Tokyo Metro. Res.Lab.P.H., 38, 423 ~ 430, 1987 (in Japanese with English abstract)

OECD SIDS	BENZOYL PEROXIDE
5. TOXICITY	ID 94-36-0
	DATE: AUGUST 2002
(c)	
Туре	: Bacterial reverse mutation assay
System of testing	: Salmonella typhimurium TA97a, TA100, TA102, TA104
Concentration	: 1 - 100 µg/plate
Metabolic	: With and Without
activation	
Results	
Cytotoxicity conc	: Not mentioned
Precipitation conc	: With metabolic activation : negative
Genotoxic effects	: Without metabolic activation : negative
	: Other (Ames, 1973) • No data
ULP Tast substance	: No dala • Denzovil nerovide Source Aldrich Chemical Co. nurity - unknown
Remarks	 Method : Preincubation method modified by Ames
ixemarks	Benzovl perovide was non-mutagenic when tested using Salmonella
	typhimurium TA97a TA100 TA102 TA104 strains with or without
	metabolic activation
Reliability	: (2) Reliable with restriction
References	: Dillon, D., Combes, R. and Zeiger, E., The Effectiveness of
	Salmonella strains TA100, TA102 and TA104 for Detecting
	Mutagenicity of Some Aldehydes and Peroxide, Mutagenesis, 13 (1),
	19 ~ 29, 1998
(d)	
Type	: Bacterial reverse mutation assay
System of testing	: Salmonella typhimurium 1A9/, 1A98, 1A100, 1A1535, 1A153/
Concentration	: A nair log dose intervals up to a dose that elicited toxicity, five doses
Matabalia activation	with and Without
Results	
Cytotoxicity conc	• Judged by a preliminary test, but no data
Precipitation conc	: With metabolic activation : negative
Genotoxic effects	• Other (Hewerth et al. 1982)
CLD	• No data
ULI Test substance	· Benzovil perovide Source - Aldrich Chemical Co purity =
Test substance	unknown)
Remarks	Method · Preincubation method
Reliability	: (4) not assignable
	Insufficient experimental details
References	: Zeiger, E., Anderson, B., Haworth, S., Lawlor, T., Mortelmans, K.
	Salmonella Mutagenicity Tests : IV. Results from the Testing of 300
	Chemicals. Env. Mol. Mutag., 11/12, 1 ~ 158, 1988

B. NON-BACTERIAL TEST

(a)				
Туре	: Mammalian Chromosomal aberration test in vitro			
System of testing	: Species/Strains : Chinese hamster cultured cell/CHL strain			
Concentration	: The max. dose 0.2 mg/ml			
Metabolic activation	: Without			
Results	:			
Cytotoxicity conc	 With metabolic activation : not done Without metabolic activation : > 0.2 mg/ml (upper limit concentration to allow the proliferation of CHL cells) With metabolic activation : 			
r recipitation conc	: with metabolic activation :			
Genotoxic effects	: Without metabolic activation : negative			
Method	: In vitro mammalian cytogenetic test (method of Ishidate, 1976)			
GLP	: No data			
Test substance	: Diluted benzoyl peroxide (used as a food additive in flour)			
Remarks	: Negative at 0.2 mg/ml, the upper limit concentration to allow the cell proliferation, with respect to frequency of polyploidy, chromosome aberration, D20 value and TR value, after 48 hours of treatment.			
Reliability	: (2) Reliable with restrictions			
	The study was undertaken as a screening test for chromosomal aberration <i>in vitro</i> . Though the documentation is not sufficient, it should be scientifically acceptable.			
References	: Ishidate, M.J., Yoshikawa, K. and Soluni, T., Mutagenicity Testing of Food Additives-First Screening Data Hegen-to-dokusei, Mutagens Toxicol., 3, 82 ~ 90, 1980			

5.6 GENETIC TOXICITY IN VIVO

(a) Preferred Result : mammalian erythrocyte micronucleus test in vivo Type Species/Strains : Mouse/ICR (SPF) Sex : Male Method : OECD TG 474 Year : 2001 GLP : Yes Route of administration : i.p. Dose : 0, 50, 100, 200 mg/kg b.w Exposure period : daily treatment for 2 days Statistical methods : Chi-square test Exposure period : single dose : 0, 50, 100, 200 mg/kg b.w Doses

OECD SIDS			BENZOYL PEROXIDE	
5. TOXICITY			ID 94-36-0	
			DATE: AUGUST 2002	
Test substance Test condition Results	 Benzoyl peroxide, s Age at study initiation No. of animals per dos Vehicle : Pluronic F-60 Duration of test : 2 day Frequency of treatmen Sampling times and treatmen Samples/dose Control groups and treatmen Clinical observations p Organs examined 	source - Sigma (B-2030), 5 : 7 weeks Se : 6 8 solution (5 %) /s tt : one daily treatment number of samples : 24 hours eatment : negative control (Plur omycin C) performed : mortality, general c <i>at necropsy</i> : <i>not examine</i>	purity = 71.8 % s after last administration, 6 conic F-68 solution), Positive linical observations od	
	Effect on mitotic ind	lex or PCE/NCE ratio by	dose level	
	Dose (mg/kg)	Group mean (PCE/PCE+NCE)	Group mean frequency of MNPCE (per 1000)	
	Vehicle	0.54	1.33	
	50	0.57	0.75	
	100	0.52	0.83	
	200	0.45	1	
	Positive control	0.43	59.7	
Genotoxic effects	 Negative statistical results : Only positive control group showed statistically increased frequency of micronucleated cells (p < 0.01). Description, severity, time of onset and duration of clinical signs : On the 2nd day, hypoactivity was observed at the concentration of 200 mg/kg. Body weight changes by dose and sex : On the 3rd day for the sacrifice, there was a decrement of body weight at the concentration of 100 and 200 mg/kg (p < 0.05). Food/water consumption changes by dose and sex : none 			
Conclusion	: Benzoyl peroxide s vivo up to the test co	showed negative result is oncentration of 200 mg/kg	n micronucleus test <i>in</i>	
Reliability	: (1) Reliable without rest This study was c guideline.	rictions onducted by GLP lab	based on OECD test	
References	: National Institute Micronucleus test tested by LGCI), 20	of Environmental Resea of Benzoyl peroxide in 001	arch (NIER), Korea ; mouse (test No.S349,	
(b)				
Туре	: A dominant lethal a	ssav		
Species/Strains	: Mouse/ICR (Ha Sw	viss)		
Sex	: Male/Female)		
Route of Administration	: i.p			
Exposure period	: 8 weeks (total week	as of mating)		
Doses	: 7 males at the lower mg/kg), 10 males for	r dose (54 mg/kg), 9 male or the control group (solve	es at the higher dose (62 ent)	
Results	: Negative			
Method	: Modified dominant	lethal mouse assay		

OECD SIDS	BENZOYL PEROXIDE
5. TOXICITY	ID 94-36-0
	DATE: AUGUST 2002
GLP	: No data
Test substance	: Source : K & K Laboratories. Inc.
Remarks	 Source : Reck Eaboratories, inc. All females were sacrificed at 13 days after mating. Each male mouse was treated with a test chemical or with the solvent alone, and subsequently caged with 3 untreated virgin female mice, which were replaced weekly for 8 Consecutive weeks. All females were sacrificed at 13 days after the midweek of their caging and presumptive mating, without being checked for vaginal plugs. Each female was scored for pregnancy, and for numbers of total implants, as comprised by live implants, early fatal deaths and late fatal deaths Total implants, early fetal deaths (brown or black containing necrotic and hemorrhagic material but no embryo), late fetal deaths (implantation sites containing a subnormal fetus and placenta which may be white or pale) were used as Criteria for evaluating results. This test was a modified dominant lethal mouse assay and benzoyl peroxide was proved to be not meeting any screening criteria for
Reliability References	 mutagenic effects. (2) Reliable with restriction Epstein, S.S., Arnold, E., Andrea, J., Bass, W. and Bishop, Y., Detection of Chemical Mutagens by the Dominant Lethal Assay in the Mouse. Toxicology and Applied Pharmacology 23, 288 ~ 325
	1972

5.7 CARCINOGENICITY

(a) Preferred Result	
Test type	: 2-year dermal carcinogenicity study
Species	: Mouse
Strain	: B6C3F1/CrlBR
Sex	: Male and female
Vehicle	: aqueous carbopol gel (0 % BP)
Method	: Covance Protocol 6711 - 100 (US FDA)
Year	: 2001
GLP	: Yes
Test substance	: Benzoyl peroxide in carbopol gel (1, 5 and 25 - 15 %)
Test Conditions	
Route of Administration	: Dermal
Exposure period	: at least 104 weeks
Frequency of treatment	: once daily
Post exposure observation period	: none
Doses	: 0, 1, 5, 25 mg/mouse/day
Control group	: Yes, vehicle control, untreated control, sentinels (not received treatment)
Area covered	: ca. 2×3 cm of dorsal skin
Occlusion	: semi-occlusive
Concentration in vehicle	: 1, 5, 15 and 25 % BP
Total volume applied	: 0.1 mL

Test conditions
Mice in Group 1 serve vehicle controls and received daily topical applications of the carbopol gel vehicle at a dose volume of 0.1 mL. Mice in Group 6 served as negative controls ; the hair on the backs of these mice was clipped at the same intervals as the other mice on study; however, these mice were not treated. Sixty mice/sex were assigned to Groups 1, 2, 3, 4, and 6, with the first 10 mice/sex/group designated for interim sacrifice during week 53 and the remaining 50 mice/sex/ group designated for terminal sacrifice after 104 weeks of treatment. Fifty mice/sex in Group 5 served as recovery animals, in that they were treated with 25 % benzoyl peroxide for 52 weeks, then treated with the vehicle for the remainder of the study. Twenty mice/sex in Group 7 served as sentinel animals for pathogen screening at Weeks 26, 52, 78, and 104.

Diet and water were provided *ad libitum*. Once weekly, each animal was removed from its cage and examined for abnormalities and signs of toxicity, specifically noting the location, size, and appearance of any grossly visible or palpable masses. The treated skin for analogous site on the untreated control) was graded for irritation once weekly. Body weights were recorded weekly from weeks 1 through 14 and every fourth week thereafter and at weeks 53 and f 5. Group 4 had additional body weights taken at weeks 57 and 59. Food consumption was measured and recorded weekly for weeks 1 through 13 and every fourth week thereafter and at weeks 52 and 104. Blood smears were prepared from all moribund-, interim-, and terminal sacrifice animals for possible evaluation of hematopoietic neoplasia.

After 52 weeks (interim-sacrifice animals) or 104 weeks (terminal sacrifice animals) of treatment the animals were anesthetized, weighed, exsanguinated, and necropsied. At necropsy, macroscopic observations were recorded, and selected tissues were collected and preserved. The liver with gallbladder, kidneys, and brain were weighed from all animals at interim sacrifice. Selected tissues (treated skin, untreated skin, and livers) were examined microscopically from all interm-sacrifice mice (Groups 1, 2, 3, 4, and 6)

All collected tissues were examined microscopically from all terminal-sacrifice mice in Groups 1, 2, 3, 4, and 6, whereas only treated and untreated skin was examined microscopically from the mice in Group 5 (high-dose-discontinued). Tumors were statistically analyzed separately and combined for relationship to treatment.

BENZOYL PEROXIDE
ID 94-36-0
DATE: AUGUST 2002

Results	: With the exception of the findings of ulcerations at the application site, there were no treatment-related differences in clinical observations among any of the groups. Treatment did not affect survival, body weights, or food consumption. The major microscopic findings were observed at the application site. At the interim (Week 53) sacrifice, these findings consisted of hyperkeratosis (1, 5, and 25 % benzoyl peroxide), subepidermal subacute inflammation (5 and 25 % benzoyl peroxide), and sebaceous gland hyperplasia (males : 5 and 25 % benzoyl peroxide ; females : 1, 5, and 25 % benzoyl peroxide). These findings were dose-dependent with regards to incidence and/or group mean severity. Similar findings were noted at the terminal (Week 105 - 106) sacrifice with the exception that there were no findings for treated or untreated skin in the high-dose and high-dose-discontinued animals. The high-dose-discontinued animals, after being treated with 25 % benzoyl peroxide for 1 year, were allowed 52 weeks of recovery.
---------	---

Although the high-dose animals were not originally intended to be a recovery group, they were not treated with benzoyl peroxide for the final 13 weeks of study since the 25 - 15 % benzoyl peroxide concentrations had been found to exceed the maximum tolerated dose. In both of these cases there was no residual effect of treatment.

- : Under the condition of this study, there were no histologic findings indicative of oncogenicity resulting from daily topical exposure of mice to benzoyl peroxide gels at concentration up to 25 % (dose level 25 mg/mouse/day). The spectrum of neoplasms observed in this study was typical for aging B6C3F1 mice.
 - : (1) Reliable without restriction
- : Consumer Healthcare Products Association (CHPA), Dermal Oncogenicity Study of Benzoyl Peroxide Gels in Mice, Covance study No.6711-100, Vol 10f 7, 2001

(b)	
Test type	: initiation/promotion
Species/Strains	: Mouse/SENCAR
Sex	: Female
Route of Administration	: Skin
Method	: Other (one year skin tumor induction test)
Year	: 1981 (paper received)
GLP	: Not mentioned
Test substance	: Benzoyl peroxide (CAS No. 94-36-0)
Test Conditions	:
Exposure period	: 53 weeks
Frequency of treatment	: 2 times a week
Post exposure observation	: none
eriod	
Doses	: 1, 10, 20, 40 mg of Benzoyl peroxide in 0.2 ml of acetone
Control group	: Yes, Concurrent vehicle

Conclusion

Reliability

References

OECD SIDS	BENZOYL PEROXIDE
5. TOXICITY	ID 94-36-0
	DATE: AUGUST 2002
Dose applied	: 1) tumor promoting activity (Group 1 - 4) : single topical application of 10 nmole of DMBA in 0.2 ml of acetone. After one week, topical applications of various dose levels of benzoyl peroxide was applied twice weekly for 52 weeks.
	 2) tumor initiating activity (Group 6 - 9) : one topical application of various dose levels of benzoyl peroxide in 0.2 ml of acetone. After one week, 2 μg of TPA was applied twice weekly for 52 weeks. 3) Complete carcinogenic activity (Group 11 - 14) : topical applications of various dose levels of benzoyl peroxide in 0.2 ml of acetone twice
	weekly for 52 weeks 4) control (Group 5, 10, 15) : acetone only, 0.2 ml twice weekly for 52 weeks
Age at study initiation	: 7 - 9 weeks
No. of animals clinical observations performed and frequency	 : 30 animals/dose : number and incidence of papillomas and carcinomas were recorded
Results	: Although benzoyl peroxide is not a complete skin carcinogen or a skin tumor initiator, it is an effective promoter of both papillomas and squamous cell carcinomas. Even at a dose of 40 mg given twice weekly for 1 year, benzoyl peroxide was not effective as a complete carcinogen. Benzoyl peroxide did not show any tumor-initiating activity when it was applied once at various dose levels, with subsequent repetitive applications of the known promoter TPA. However, when benzoyl peroxide was applied topically twice weekly after initiation of tumors with DMBA, it proved to be an effective promoter with a reasonable dose-response relationship. As little as 1 mg given twice weekly produced a significant number of papillomas and carcinomas. The tumor response appeared to plateu at a dose level of 20 mg, with 79 percent of the mice developing papillomas (average of 5.2 papillomas per mouse at 30 weeks). At the same dose level, 43 percent of the mice developed squamous cell carcinomas at 52 weeks.

induction	duction of bezoyl peroxide in SENCAR mice					
	Group	Dose (mg)	Mice Alive at 30 weeks	Papillomas/ mouse at 30 weeks	Mice with papillomas at 30 weeks (%)	Mice with carcinomas at 52 weeks (%)
	Tumor p	promoting activi	ty (after DM	BA initiation)	
	1	1	29	0.8	32	4
	2	10	28	3.8	72	22
	3	20	27	5.2	79	43
	4	40	24	5.4	85	40
	5	Acetone (only 0.2ml)	28	0.03	3	0
	Tumor i	nitiating activity	/ (with TPA	promotion)		
	6	1	29	0.1	10	0
	7	10	28	0.1	10	0
	8	20	28	0.1	10	0
	9	40	27	0.2	15	0
	10	Acetone (only 0.2ml)	29	0.2	15	0
	Comple	te carcinogenic a	activity			
	11	1	28	0	0	0
	12	10	29	0.03	3	0
	13	20	27	0.03	3	0
	14	40	25	0.03	3	0
	15	Acetone (only 0.2ml)	29	0	0	0
		1 . 1	. 11	.1 .11		1 .

Skin tumor

Benzoyl peroxide promoted both papillomas and carcinomas when it was topically applied to mice after DMBA initiation. Benzoyl peroxide was inactive on the skin as a complete carcinogen or as a tumor initiator.

- : (2) Reliable with restriction
- : Slaga, T.J., Klein-Szanto, A.J.P., Triplett, L.L.I., Yotti, L.P. and Trosko, J.E., Skin Tumor-Promoting Activity of Benzoyl Peroxide, a Widely Used Free Radical-Generating Compound, Science, 213, 1023 ~ 1025, 1981

(c)	
Species/Strains	: Rat and Mouse/Albino
Sex	: Male/Female
Route of Administration	: Oral intake, subcutaneous injection, Dermal painting
Method	
GLP	: No data
Exposure period	: Oral - 120 weeks in rat, 80 weeks in mouse ; Subcutaneous-120 weeks in rat, 80 Weeks in mouse ; Dermal-80 weeks in mouse

Reliability

References

OECD SIDS	BENZOYL PEROXIDE
5. TOXICITY	ID 94-36-0
	DATE: AUGUST 2002
Frequency of treatment	: Oral - ad libitum ; Subcutaneous-single injection/week; Dermal-6 times a week
Post exposure observation period	: Not mentioned
Doses	: Oral - 28, 280, 2,800 mg/kg diet ; Subcutaneous - 120 mg/rat, 50 mg/mouse : 50 mg (50 % suspension)
Control group	: Yes Concurrent no treatment Concurrent vehicle
Test substance	: Benzovl peroxide purity = unknown
Results	: The overall tumor incidence was not significantly different between the experimental and control groups in rats or mice
Reliablity	: (2) Reliable with restriction
References	: Sharratt, M., Frazer, A.C. and Forbes, O.C., Study of the Biological Effects of Benzoyl Peroxide, Food Cosmet. Toxicol., 2, 527 ~ 538, 1964
(d)	
Species/Strains	: Mouse/SENCAR
Sex	: Female
Route of Administration	: Dermal
Method	:
GLP	: No data
Exposure period	: Dermal - 100 weeks
Frequency of treatment	2 times a week
Post exposure observation period	: not mentioned
Doses	: 0.1, 0.3, 0.6, 2.0 and 20 mg
Control group	: No data
Test substance	: Benzoyl peroxide, purity = unknown
Results	: Benzoyl peroxide is a mouse skin promoter but was inactive as a complete carcinogen or as a tumor initiator.
Reliability	: (2) Reliable with restriction
Keferences	: Gimenez-Conti, I.B., Oubari, D., and Slaga, T.J., Dose-response of benzoyl peroxide hyperplastic promoter and possible complete carcinogenic activities, Proc. Am. Assoc., Cancer Res., 34, 184, 1993
(e)	
Species/Strains	• Mouse/hr/hr Oslo strain and SENCAR
Sex	 Moles/m/m Oslo strain and SERVERIK Male/Female
Route of Administration	• Skin
Method	• Skii
GLP	• • No data
Exposure period	• 52 Weeks
Frequency of treatment	: Twice a week
Post exposure observation	: none
period	• 10110
Doses	: 5 % Benzoyl peroxide in a gel (Panoxyl)
Control group	: Yes
	Concurrent vehicle, Historical

OECD SIDS	BENZOYL PEROXIDE
5. TOXICITY	ID 94-36-0
	DATE: AUGUST 2002
Test substance	· Banzovi parovida purity - unknown
	· Denzoyi peroxide, purity – unknown
Results	: No induction of carcinoma
Reliability	: (2) Reliable with restriction
References	: Iversen, O.H., Skin Tumorigenesis and Carcinogenesis Studies with
	7,12-dimethylbenzene[a]anthracene, ultraviolet Light, Benzoyl
	Peroxide (Panoxy gel 5 %) and Ointment Gel Carcinogenesis, 9 (5), 803 ~ 809, 1988
(f)	
Species/Strains	: Mouse/SENCAR
Sex	: Female
Route of Administration	: Skin
Method	
GLP	: No data
Exposure period	: 20 Weeks
Frequency of treatment	: 2 times a week
Post exposure observation	: None
period	
Doses	: 20 mg of Benzoyl peroxide in 0.2 ml of acetone
Control group	: Yes
Test substance	: Benzoyl peroxide, purity = unknown
Results	: Benzoyl peroxide enhanced the progression of preexisting
D 11 1 11	papillomas.
Reliability	: (2) Reliable with restriction
References	: O'Connell, J.F., Klein-Szanto, A. J.P., DiGiovanni, D.M., Fries,
	J.W. and Slaga, T.J., Enhanced Malignant Progression of Mouse
	Skin Tumors by the Free-Radical Generator Benzoyl Peroxide,
	Cancer Research, 46 (6), $2863 \sim 2865$, 1986
(g)	
Species/Strains	: Mouse/Uscd (Hr) stock albino
Sex	: No data
Route of Administration	: Skin
Method	:
GLP	: No data
Exposure period	: 70 Weeks
Frequency of treatment	: 5 times a week
Post exposure observation	: none
Doses	• 5 % Panzoul paravida lation in an aquaous vahiala
Control group	• Vas Concurrent vehicle
Test substance	 Ponzovi porovido, purity = unknown
Results	: Denzovi peroxide, pullty – ulikilowii : Ponzovi peroxide, did not set as skin tumor promotors under this
Results	andition in Which croton oil promoted skin tumors following
	initiation by LIV radiation
Reliability	• (2) Reliable with restriction
References	• Enstein I Photocarcinogenesis Promotion Studies with Renzovi
	Peroxide and Croton Oil 1 Investigative Dermatology 91(2) 114~
	116, 1988

OECD SIDS	BENZOYL PEROXIDE
5. TOXICITY	ID 94-36-0
	DATE: AUGUST 2002
(h)	
(II) Species/Strains	• Hamster/Pathagan free Surian golden
Sev	• Mala
Boute of Administration	
Method	• SKIII
GLP	• • No data
Exposure period	• 64 wooks
Exposure period	• 04 weeks
Post exposure observation	· 5 times a week
period	•
Doses	: 80, 160 mg of benzovl peroxide in 1 ml of acetone
Control group	: Yes. Concurrent vehicle
Test substance	: Benzovl peroxide, purity = unknown
Results	: Benzovl peroxide is able to promote the formation of dermally
	located Melanotic tumors
Reliability	: (2) Reliable with restriction
References	: Schweizer J Loehrke H Edler L and Goerttler K Benzovl
	Peroxide promotes the Formation of Melanotic Tumors in the Skin
	of 7 12-dimethylbenz[a]anthracene -initiated Syrian Golden
	Hamsters Carcinogenesis $8(3)$ 479 ~ 482 1987
	Tumsters, Caremogenesis, 6 (3), 173 162, 1967
(i)	
Species/Strains	: Hamster/Syrian golden
Sex	: Male/Female
Route of Administration	: Skin (painting of the hamster buccal pouch)
Method	
GLP	: No data
Exposure period	: 6 weeks
Frequency of treatment	: 3 times a week
Post exposure observation	: none
period	
Doses	: 40 % solution of Benzoyl peroxide in acetone (equivalent to 20 mg
	of benzoyl Peroxide per single brush application)
Control group	: Yes
	Concurrent no treatment, Concurrent vehicle
Test substance	: Benzoyl peroxide, purity = unknown
Results	: Benzoyl peroxide is noncarcinogenic, but it in a 40 % solution of
	benzoyl peroxide in acetone acts as a promoter after initiation with
	a subthreshold dose of 0.1 % DMBA in oil.
Reliability	: (2) Reliable with restriction
References	: Odukoya, O. and Shklar, G., Initiation and Promotion in
	Experimental Oral Carcinogenesis, Oral Surg. Oral Med. Oral
	Pathol., 58 (3), 315 ~ 320, 1984
(i)	
U) Species/Strains	· Mouro/SENCAD
Species/Sitallis Sev	· IVIOUSC/SEINCAK
Boute of Administration	: remaie
Method	• 5KIII
GLP	• • No data
	• IN uala

OECD SIDS	BENZOYL PEROXIDE
5. TOXICITY	ID 94-36-0
	DATE: AUGUST 2002
Exposure period	: 51 weeks
Frequency of treatment	: 2 times a week
Post exposure observation	: none
period	10 ma of harmoul narrouids in 0.2 ml of agatana (against to 2ma
Doses	honzovi Derovido)
Control group	Vez Company trachicle
Test substance	Personal nervide numity - university
P equite	Benzovi peroxide, purity – unknown
Results	arcinogonia action
Reliability	(2) Poliable with restriction
References	· (2) Kellable with restriction
References	Havashi V. Studios on the Dramating and Complete Carringgonia
	Activities of Some Oxidizing Chemicals in Skin Careinogenesis
	Concor Letters 24, 200 204, 1084
	Cancer Letters 24, 29978 304, 1984
(k)	
Species/Strains	: Mouse/Swiss Albino
Sex	: Female
Route of Administration	: Skin
Method	:
GLP	: No data
Exposure period	: 46 weeks
Frequency of treatment	: 2 times a week
Post exposure observation	:
period	
Doses	: $20 \text{ mg}/300 \mu\text{I}$
Control group	: Yes
Tast - hatawa	Concurrent no treatment
l est substance	: Benzoyl peroxide, purity = unknown
Results	: Iron overload augments Benzoyl peroxide-mediated tumor
	promotion in 7,12-dimethylbenzen[a]anthracene (DMBA)-initiated
Paliability	mouse skin.
References	: (2) Kellable with restriction • Degradaby II and Athen M. Effect of Iron Overload on the
Kelefences	Rezazadell, H., and Athal, M., Effect of fion Overload of the
	Cancer Letters 126, 135 - 142, 1008
	Cancer Letters, 120, $155 \approx 142$, 1998
(1)	
Sex	: Male/Female
Route of Administration	: Dermal
Method	
GLP	: No data
Exposure period	: 164 days in Car-S (Carcinogenesis-susceptible) strain, 253 days in
-	Car-R (Carcinogenesis-resistant) strain
Frequency of treatment	: 2 times a week
Post exposure observation	: Promotion by exposure with chemicals was continued as long as the
period	increase of papilloma number was a linear function of time, at least
	in the most susceptible line.
Doses	: 3.7, 7.5 and 15.0 mg Benzoyl peroxide

OECD SIDS	BENZOYL PEROXIDE
5. TOXICITY	ID 94-36-0
	DATE: AUGUST 2002
Control group	. V
Tost substance	
Test substance	: Benzoyl peroxide, purity – unknown Mauga/Car D. and Car S. mias meducad by 16 consecutive
Test condition	generation of bi-directional selective breeding at the ENEA (Rome, Italy)
Results	: Benzoyl peroxide is also able to promote carcinoma development in Car-S mice
Reliability	: (2) Reliable with restriction
References	: Saran, A., Pazzaglia, S., Rebessi, S., Bouthillier, Y., Pioli, C., Covelli, V., Moutoin, D., Doria, G. and Biozzi, G., Skin Tumorigenesis by Initiators and Promoters of Different Chemical Structures in Lines of Mice Selectively Bred for Resistance (Car-R) or Susceptibility (Car-S) to Two-stage Skin Carcinogenesis, Int. J. Cancer, 83 (3), 335 ~ 340, 1999
(m)	
Test type	: Tumor promotion using human bronchial epithelial cell culture
Species/Strains	: Normal human bronchial epithelial cell
Method	:
GLP	: No data
Exposure period	: 6 hours exposed to Benzoyl peroxide and followed by 7 - 9 days of incubation for the tests of cell proliferation and morphological changes; one hour for DNA damage test.
Frequency of dosing	: single dosing
Number if replicates	: duplicate dishes or well were measured and each experiment was repeated 3 times for the tests of cell proliferation and morphological changes.
Doses	: 0.003, 0.03 and 0.3 mM in cell proliferation test ; 0.056 mM in morphological Change test ; 0.1 mM in DNA damage test
Control group	: Yes Concurrent no treatment
Test substance	: Benzoyl peroxide, purity = unknown
Results	: The 50 % of control concentration (growth rate) for the 6-hr exposure was found to be 0.065 mM Benzoyl peroxide ; Benzoyl peroxide–exposed cells were smaller than controls (median cell planar area, 620 sq μm versus 1,150 sq μm) ; Benzoyl peroxide caused detectable amounts of both single strand breaks and DNA-protein cross-links.
Reliability	: (2) Reliable with restriction
References	: Saladino, A.J., Willey, J.C., Lechner, J.F., Grafstrom, R.C., LaVeck, M. and Harris, C.C., Effects of Formaldehyde, Acetaldehyde, Benzoyl Peroxide, and Hydrogen Peroxide on Cultured Normal Human Bronchial Epithelial Cells, Cancer Research, 45, 2522 ~ 2526, 1985

5.8 TOXICITY TO REPRODUCTION

OECD SIDS		BENZOYL PEROXIDE
5. TOXICITY		ID 94-36-0
		DATE: AUGUST 2002
(a) Preferred Result		
Species/strains	:	Rat/Sprague-Dawley
Sex	:	Male/Female
Method	:	Combined Repeated Dose and Reproduction/Developmental Toxicity Screening Test OECD TG 422
Year	:	2001
GLP	:	Yes
Route of administration	:	Oral (gavage)
Dose level	:	0, 250, 500, 1,000 mg/kg b.w/day
Exposure period		Male : 29 days, female : from 2 weeks before mating to the day 3 of lactation
Frequency of treatment	:	Daily
Control group	:	(-) Concurrent vehicle (+) Cyclophosphamide (4.5 mg/kg/day)
Post exposure	:	1 dav
observation period	•	
Premating exposure period	:	14 days
Statistical methods	:	Dunnett's multiple comparison test
Test substance	•	Source · SIGMA purity · 70 %
Test condition	•	- Test Subjects Age · 7 week old for male and female
	•	- Weight at study initiation : 323 38 - 356 63 g for males 212 21 -
		237.18σ for females
		- No of animals : 10 animals/sev/dose (5 animals/sev/positive
		control)
		- Vehicle : Corn oil
Frequency of treatment Control group Post exposure observation period Premating exposure period Statistical methods Test substance Test condition	:::::::::::::::::::::::::::::::::::::::	Daily (-) Concurrent vehicle (+) Cyclophosphamide (4.5 mg/kg/day) 1 day 14 days Dunnett's multiple comparison test Source : SIGMA, purity : 70 % - Test Subjects Age : 7 week old for male and female - Weight at study initiation : 323.38 - 356.63 g for males, 212.21 - 237.18 g for females - No. of animals : 10 animals/sex/dose (5 animals/sex/positive control) - Vehicle : Corn oil

- Dosing schedule : males – premating period (2 weeks) and mating period ; 29 days females-premating period (2 weeks), mating, pregnant and lactation period ; $41 \sim 51$ days

- Mating procedures : M/F ratios per cage ; 1/1, proof of pregnancy ; sperm detection in vagina

- Clinical observations performed and frequency : General condition was observed once a day. Body weight and food consumption of pregnant animals were determined on the day 1, 7, 14, 20 of pregnancy and the day 1, 3 of lactation. Haematology and biochemistry for males only at the time of necropsy after 28 days of chemical exposure

- Sperm examination : concentration, motility, morphology was observed at necropsy.

- Organs examined at necropsy (macroscopic and microscopic) : Organ weight : liver, kidney, adrenal gland, testis, epididymis, thymus, spleen, brain, heart, ovary, uterus, thyroid gland, spermary, prostrate gland. Microscopic : brain, spinal cord, stomach, pancreas, jejunum, ileum, cecum, colon, rectum, liver, kidney, adrenal gland, spleen, heart, thymus, thyroid, bronchus, lung, pituitary gland, ovary, uterus, vagina, testis, epididymis, spermary, prostrate gland, mammary gland, bladder, nodi lymphatici mesenterici, nodi lymphatici, nervus ischiadicus, femoral marrow and all gross lesion. - Parameters assessed during study P and F1 : copulation index (matings/pairs x 100), fertility index (Pregnancies/matings x 100), gestation index (live litters/pregnancies x 100), viability index (No. of live offspring at day1 or day3/No. of live offspring at birth \times 100), body weight of live pups (on day 0 and 4), No. of corpora lutea, No. of female mated, abortion, premature birth, gestation period, sex ratio (Total No. of male pups/Total No. of female pups)

NOAEL parental NOAEL F1 Offspring NOAEL F2 Offspring Results

- = 500 mg/kg/day
- : = 500 mg/kg/day
- := N/A
- : Significant decrease in absolute and relative weights of left testes and relative wt of epididymis was observed in male animals exposed to 1000 mg/kg (p < 0.05). Severe testis atrophy was seen in the highest exposure male group but these changes were not seen in recovery animals. The pups which exposure to highest levels of benzoyl peroxide showed significant decrease in body wt. Gain (p < 0.01). There was no significant difference between experimental and control groups in the precoital time, copulation, fertility and gestation rate.

DOSE : (mg/kg)	0	250	500	1000	Cyclophos phamide
No. of mated males	10	10	10	10	10
Copulation index (%)	100	100	100	100	100 (5/5)
Fertility index (%)	100	100	100	90 (9/10)	100 (5/5)
No. of mated females	10	10	10	10	10
Copulation index (%)	100	100	100	100	100 (5/5)
Fertility index (%)	100	100	100	90 (9/10)	100 (5/5)
Gestation index (%)	100	100	100	100	40 (2/5)
No. of dams	10	10	10	9	2
No. of corpora lutea	169	182	171	157	90
Mean±S.D	16.9±4.1	18.2±2.3	17.1±1.9	17.4±2.5	18±1.9
No. of implantations	150	150	161	136	81
Mean±S.D	15.0±1.8	15.0±2.9	16.1±1.9	15.1±2.1	16.2±1.6
Mean % preimplantation loss	9	16.1	5.8	12.9	9.9
No. of embryo/fetal death	10/3	9/3	7/3	3/0	64/9
No. of live pups born					
Mean±S.D	13.7±2.7	13.8±2.5	15.1±2.1	14.8±2.2	1.6±2.3*
Mean pregnancy period (day)	20.4	20.8	20.6	20.8	22
Viability index on day 1 pp	97.5	98.1	100	100	53.3*
Viability index on day 3 pp	100	100	100	100	90*
Body weights of pups (g)					
Male 1 DAY	6.88±0.53	7.22±0.81	6.91±0.64	6.72±0.79	4.59±0.77
3 DAY	8.93±0.91	9.10±1.03	8.69±0.79	8.12±1.01*	7.16±0.88
Female1 DAY	6.51±0.53	6.93±0.76*	6.73±0.59	6.21±0.79	4.33±0.74
3 DAY	8.60±0.99	8.79±1.03	8.48±0.89	7.49±1.03*	5.48±0.89
Precoital time					
Mean±S.D (day)	2.2±1.2	3.2±1.3	2.4±1.2	2.3±1.2	5.3±5.3

- Mortality and day of death : No deaths were found in all animals including control groups.

- Body weight : Body weight gains were slightly increase (approximately 7 % more than controls) in female recovery group exposed to 1,000 mg/kg only after 29 days. No significant body weight changes were observed in all treatment animals except highest dose recovery group. Low body weight gain of pups showed in 1,000 mg/kg dose.

- Food/water consumption : The mean food consumption of 1,000 mg/kg was significantly increased at first and fourth weeks of chemical exposure in male. No other treatment groups were showed abnormal food consumption.

- Organ weight changes : Significant decrease in absolute and relative weights of left testes and relative wt of epididymis were observed in 1000 mg/kg exposure group, however these changes were not seen in recovery group.

- Histopathology incidence and severity : In the 1,000 mg/kg treatment group, testis degenerations were observed microscopically and these pathosis were mainly morphological change in sperm cell such as apoptosis, cellular swelling and multinucleated giant cell. In the female reproductive organ, slight affects were observed in uterus exposed to 1,000 mg/kg.

- Grossly visible abnormalities, external, soft tissue and skeletal abnormalities.

Treatment (mg/kg/day)	0	250	500	1000	Cyclophosphamide
No. with major abnormalities	0	1	0	0	0
Mean % of pups examined	0		0	0	0
		0.6			
No. of females affected	0	1	0	0	0
No. with minor abnormalities	20	20	18	60	8
Mean % of pups examined	14.	12.	11.		100.0*
	2	4	5	41.4*	
No. of females affected	6	4	5	6	2
No. with variants	0	0	0	0	0

In the 1000 mg/kg dose group, the incidence of runts was significantly higher then control group.

* Statistically significant difference from control group (p < 0.01)

Conclusion

: Significant toxic effect was found in P0 and F1 exposed to 1,000 mg/kg. This result indicated that the NOAEL for both reproduction and developmental toxicity was 500 mg/kg b.w/day.

Reliability

: (1) Reliable without restriction

References : National Institute of Environmental Research (NIER), Combined Repeated Dose Toxicity with the Reproduction/Developmental Toxicity Screening Testing of Benzoyl peroxide in Rats (Report No.P049, tested by LGCI), 2001

5.9 DEVELOPMENTAL TOXICITY/TERATOGENICITY

(a) Preferred Result				
Species/strains		Rat/Sprague-Dawley		
Sex :		Male/Female		
Route of administration	:	Oral (gavage)		
Method	:	Combined Repeated Dose and Reproduction/Developmental Toxicity Screening Test OECD TG 422		
Year	:	2001		
GLP	:	Yes		
Dose level	:	0, 250, 500, 1,000 mg/kg b.w/day		
Exposure period	:	Male : 29 days, female : from 2 weeks before mating to the day 3 of lactation		
Frequency of treatment	:	Daily		
Control group	:	(-) Concurrent vehicle (+) Cyclophosphamide (4.5 mg/kg/day)		
Post exposure	:	1 day		
observation period				
Statistical methods	:	Dunnett's multiple comparison test		
NOAEL for	:	= 500 mg/kg/day		
developmental				
Test substance	:	Benzoyl peroxide, Source-SIGMA, purity = 70%		
Results	:	The body weight gain of pups on the day 3 after birth decreased at 1,000 mg/kg. Benzoyl peroxide has adverse effects on development of pups with high birthrate of runt at 1000 mg/kg dose level. No significant differences appeared between the treatment and control group in any other observation.		
Reliability	:	(1) Reliable without restriction		
References	:	National Institute of Environmental Research (NIER), Combined Repeated Dose Toxicity with the Reproduction/Developmental Toxicity Screening Testing of Benzoyl peroxide in Rats (Report No.P049, tested by LGCI), 2001		

5.10 OTHER RELEVANT INFORMATION

- A. Specific toxicities No data
- B. Toxicodynamics, toxicokinetics

OECD SIDS	BENZOYL PEROXIDE
5. TOXICITY	ID 94-36-0 DATE: ALICUST 2002
	DATE. AUGUST 2002
(a) Tama	
Type Remarks	 Metabolism The transepidermal penetration and metabolic disposition of ¹⁴C-benzoyl peroxide were assessed <i>in vitro</i> (excised human skin) and <i>in vivo</i> (rhesus monkey). <i>In vitro</i>, the benzoyl peroxide penetrated into the skin, through the stratum corneum or the follicular openings, or both, and was recovered on the dermal side as benzoic acid. <i>In vivo</i>, benzoic acid was recovered from urine in amounts equivalent to 45 % and 98 % of the radiolabel following topical and i.m. administration, respectively, of ¹⁴C-benzoyl peroxide. Authors concluded that benzoyl peroxide penetrates as such into the skin layers and is converted therein to benzoic acid, which, in turn is absorbed into the systemic circulation. Renal clearance of the metabolite is sufficiently rapid as to preclude its hepatic conjugation with glycine, since following topical administration to rhesus monkeys, no hippuric acid was found in the urine, as could have been expected had a significant amount of benzoic acid passed through the liver.
Reliability References	 (2) Reliable with restriction Nacht, S., Yeung, D., Joseph, N., Beasley, J. N. Jr., Anjo, M. D. and Maibach, H.I., Benzoyl Peroxide : Percutaneous Penetration and Metabolic Disposition, <i>J. Am.</i> Acad. Dermatol., 4 (1), 31 ~ 37, 1981
(b) Type	: Metabolism
Remarks	: The effect of drug concentrations of 2.5, 5 and 10 % upon the transepidermal penetration of ¹⁴ C-benzoyl peroxide in a lotion vehicle was assessed in excised human skin and <i>in vivo</i> in the rhesus monkey. <i>In vitro</i> , penetration of benzoyl peroxide was dose-dependent, both as to rate and to amount, as measured by hourly recovery of the metabolite, benzoic acid, from the dermal side of the model. <i>In vivo</i> , the higher the concentration of benzoyl peroxide absorbed was excreted rapidly in urine as benzoic acid; no hippuric acid was detected at any time. Authors concluded that the transepidermal delivery of benzoyl peroxide, but not its metabolic disposition, is concentration-dependent, and the renal clearance of the systemically absorbed drug is so rapid that it precludes passage through the liver – therefore, no systemic toxicity due to the substance accumulation can be expected.
Reliability References	 (2) Reliable with restriction Yeung, D., Nacht, S., Bucks, D. and Maibach, H. I., Benzoyl Peroxide : Percutaneous Penetration and Metabolic Disposition. II. Effect of Concentration, J. Am. Acad. Dermatol., 9, 920 ~ 924, 1983

OECD SIDS	BENZOYL PEROXIDE
5. TOXICITY	ID 94-36-0
	DATE: AUGUST 2002
(c)	
Туре	: Metabolism
Remarks	: The greatest amount of penetrating BP (9 to 14 % of the applied dose) was found in the horny layer, which forms a reservoir and where biotransformations were reduced. Small quantities of BP diffused toward the epidermis down to the deeper dermis, where benzoic acid represented 74 % of the radioactivity. Distribution gradients of the radioactivity and conversion rates to benzoic acid were stable between 3 and 24 hours of application in all parts of the skin. This result showed that diffusion of BP from the vehicle was in balance with the dissociation processes and the blood resorption as benzoic acid throughout the experiment.
Reliability	(2) Reliable with restriction
References	: Wepierre, J., Corroller, M. and Didry, J. R., Distribution and Dissociation of Benzoyl Peroxide in Cutaneous Tissue after Application on Skin in the Hairless Rat, International Journal of Cosmetic Science 8, 97 ~ 104, 1986
(d)	
Туре	: Metabolism
Remarks	: The percutaneous penetration and the metabolism of benzoyl peroxide (BPO) were assessed <i>in vitro</i> on human skin and <i>in vivo</i> on 5 patients with leg ulcers. The BPO <i>in vitro</i> absorbed by the skin was converted to benzoic acid preferably in the dermis. The portion penetrated through the skin was benzoic acid only. Also in patients treated with BPO, no BPO could be detected in the serum. These findings show that BPO as such is absorbed by the skin, but is systemically absorbed only after the metabolisation to benzoic acid. Therefore, a systemic-toxic effect in local therapy with BPO can be excluded.
Reliability References	 (2) Reliable with restriction Morsches, B. and Holzmann, H., Untersuchungen zur perkutanen Resorption von Benzoyl peroxid, ArzneimForsch./Drug Res., 32 (1), Nr.3, 1982

5.11 EXPERIENCE WITH HUMAN EXPOSURE

(a)		
Туре	:	Epidemiological case-control study (using medical records)
Results	:	There is no evidence that acne suffers (who may have used
		benzoyl peroxide without prescription) or individuals prescribed
		benzoyl peroxide, have an increased risk of malignant melanoma.
Reliability		(2) Reliable with restriction
References	:	Cartwright, R, A., Hughes, B.R. and Cunliffe, W.J., Malignant
		Melanoma, Benzoyl Peroxide and Acne : A Pilot Epidemiological
		Case-Control Investigation, British J, Derm., 118, 239 ~ 242, 1988

OECD SIDS	BENZOYL PEROXIDE
5. TOXICITY	ID 94-36-0
	DATE: AUGUST 2002
(b)	
Туре	: Epidemiological case-control study (using cancer registry)
Results	: No statistically significant association between the prescribed use of benzoyl peroxide and skin cancer was found in 931 cases and 7630 controls.
Remarks	: Limitations were that the nonprescription use of benzoyl peroxide by cases And Controls was not considered, and the Drug Prescription Plan records did not cover the full period during which benzoyl peroxide containing medications were available.
Reliability	(2) Reliable with restriction
References	: To. T, Hogan D, Wilson, E, Miller, A, Benzoyl peroxide and facial skin cancer, Am. J. Epidemiol., 134:772
(c)	
Type	: Epidemiological case-control study (using cancer registry)
Results	: No statistically significant association was found between the use of acne medications containing benzoyl peroxide and skin cancer of the head and neck in 964 cases and 3856 controls. Also acne was not associated with increased risk of skin cancer.
Remarks	: This study covered all prescription and nonprescription use of benzoyl peroxide containing acne medications.
Reliability	(2) Reliable with restriction
References	: Hogan, D.J., To T, Wilson, E.R., Miller, A.B., Robson, D., Holfeld, K., Lane, P., A study of acne treatments as risk factors for skin cancer of the head and neck. Brit. J. Dermatol., 125, 343 ~ 348
(d)	
Type	: Contact dermatitis
Results	 Two cases were reported of allergic contact dermatitis to acne preparations containing benzoyl peroxide. Both patients had positive patch tests to 5 % Benzoyl peroxide in petrolatum while 40 of 41 control patients had no reaction.
Reliability	(2) Reliable with restriction
References	: Eaglstein, W.H., Allergic Contact Dermatitis to Benzoyl Peroxide, Arch. Derm., 97, p 527, 1968
(e)	
Туре	: Skin irritation
Results	: Two cases were reported in which topical application of benzoyl peroxide resulted in an unusual pattern of reticulate hyperpigmentation of the skin, most likely as a sequela of an irritant contact dermatitis.
Reliability References	 (2) Reliable with restriction Weinberg, J.M., Moss, T., Gupta, S.M., White, S.M. and Don, P.C., Reticulate Hyperpigmentation of the Skin After Topical Application of Benzoyl Peroxide, Acta Derm. Venereol (Stockh) 78, 301 ~ 302, 1998

OECD SIDS	BENZOYL PEROXIDE			
5. TOXICITY	ID 94-36-0			
	DATE: AUGUST 2002			
(f)				
Туре	: Skin allergy			
Results	 After 4 months of use, a patient with acne developed allergic contact urticaria to 5 % topical benzoyl peroxide. The observed symptoms were the increased hivelike lesions occurred about 20 - 30 minutes after topical application. 			
Reliability	(2) Reliable with restriction			
References	 1 kach, J.R., Allergic Contact Urticaria to Benzoyl Peroxide, Cutis 29, 187 ~ 188, 1982 			
(g)				
Type	: Allergic contact dermatitis			
Results	: Benzoyl peroxide is an irritant but has only infrequently been reported as a sensitizer. Non-medical exposures include its use as a component of acrylates and as a bleach and preservative for flour and oil. As such it has caused allergic contact dermatitis in patients in whom it was present in an acrylic bone cement and in dental acrylates. In the occupational setting it has produced allergic contact dermatitis in pakers			
Reliability	(2) Reliable with restriction			
References	 Mosby Year Book, Contact and Occupational Dermatology, St. Louis, MO/Marks, J.G., V.A. DeLeo, 164 ~ 164, 1992 			
(h)				
Туре	: Sensitization			
Results	: The investigation of patients with leg ulcers, treated with a concentrated benzoyl peroxide solution, revealed a sensitization to the substance in nearly half of the probands. Therefore and because of the probable promotion of neoplastic epidermal proliferation, benzoyl peroxide-containing solutions			
Reliability References	 (2) Reliable with restriction Von Bahmer F.A., A. Schulze-Dirks und H.Zaun, Sensibilisierende Wirkung einer fur die behandlung des Ulcus cruris verwendeten 20 % igen Benzoylperoxid-Zubereitung, Dermatosen, 32, Nr 1, 21 ~ 24, 1984 			
(i)				
Туре	: Skin irritation			
Results	: 63 patients with papulopustular acne had been treated by mean of 10 % benzoyl peroxide gel daily twice for 8 weeks at times of little insolation. Only in a few patients transient redness, burning, dryness and scaling occurred. Allergic sensitization was not observed.			
Reliability References	 (2) Reliable with restriction Haustein, U.F., Kramer, I.E., Fickweiler, Beahandlung der Akne vulgaris mittels Benzoylperoxid, Dermatol. Monatsschr., 167, 149 ~ 153, 1981 			

OECD SIDS	BENZOYL PEROXIDE
5. TOXICITY	ID 94-36-0
	DATE: AUGUST 2002
(j)	
Туре	: Patch test
Results	: A 23 years old woman with cystic acne was treated with a 5 % topical solution (Peroxiben 5 % gel). After 7 days she noted intolerance with redness and scaling and severe itching. She was patch tested with a standard series, benzoyl peroxide 1 % pet., vehicle and wood tars. The result showed a strong positive response. Another recent preparation of 1 % benzoyl peroxide in neutral gel was also positive.
Remarks	: There have been many attempts to explain contact allergy to benzoyl peroxide in patients with acne is rather uncommon compare to its capacity to sensitize. The pilosebaceous unit may generate an intermediate metabolite, the allergen being a metabolite seldom formed in facial skin, or perhaps degrade the drug more rapidly. Sweating could dilute the compound and wash away the product. Treatment without occlusive techniques would minimize the likelihood of sensitization. The presence of inflammation in acne could interfere with the induction of an allergic response. Several other reports have suggested that in many cases, the symptoms of sensitization are mild and they may be regarded merely as irritants; thus a patch test is not performed.
Reliability References	 (2) Reliable with restriction Morillas, I.M., Martinesz, A.A, J.L.S. Lozano and A.G.Perez, Is benzoyl peroxide an irritant or sensitizer?, Contact Dermatitis, 16, 232 ~ 233,1987
(k)	
Type Results	 Patch test A 25 years old man had been an electrician for the last 4 years. Benzoyl peroxide may have acted as a contactant allergen after plastic particles became airborne when sawing plastic material, or perhaps after being melted during soldering. A positive patch test to benzoyl peroxide was obtained at the recommended concentration of 1 %.
Remarks	: Allergic contact dermatitis due to benzoyl peroxide may develop as a consequence of occupational exposure, affecting mainly workers in the electronics and plastics (epoxy resins and catalysts) industries, electricians, dentists, bakers, and laboratory technicians. Airborne allergic contact dermatitis from this agent has also been reported due to molten candle wax. When benzoyl peroxide is used for therapeutic purposes, leg ulcers are more prone to develop sensitization than acneic skin.
Reliability References	 (2) Reliable with restriction Santiago, Q., J.M., Olaguibel, B.E., Garcia and Tabar, A.I., Occupational airborne contact dermatitis due to benzoyl peroxide, Contact Dermatitis. 29, 165 ~ 166, 1993

OECD SIDS	BENZOYL PEROXIDE
5. TOXICITY	ID 94-36-0
	DATE: AUGUST 2002
(1)	
Туре	: Patch test
Results	: Positive
Remarks	 272 children up to the age of 14 years were patch tested during 10 years period. An unusual feature was the finding of 10 patients patch test positive to benzoyl peroxide 1 % pet. The positivities to this allergen were classified as relevant.
Reliability	: (2) Reliable with restriction
References	dermatitis in children, Contact Dermatitis, 30, 292 ~ 294, 1994
(m)	
Type	: Patch test
Results	: 791 patients including 59 dental technician were tested with the denture material series. Benzoyl peroxide ranked 2 nd in patch test positivity. The result in dental technicians showed that positive reaction to benzoyl peroxide were observed in 9.8 % of patients. Positive and questionable reactions to benzoyl peroxide were observed in 37 and 57 other patients respectively.
Remarks	: Reactions were more frequent in dental technicians, who might be exposed to benzoyl peroxide in working environment.
Reliability	: (2) Reliable with restriction
References	: Gebhardt, M., Geier, J., Evaluation of patch test results with denture material series, Contact Dermatitis, 34, 191 ~ 195, 1996
(n)	
Type	: Patch test
Results	: Positive
Remarks	: It is reported that a 62 year-old man with total hip prosthesis developed a recurrent sterile fistula and loosening of his prosthesis due to allergy to the bone cement component benzoyl peroxide.
Reliability	: (2) Reliable with restriction
References	: Jäger, M. and Balda, B.R., Loosening of a Total Hip Prosthesis at Contact Allergy Due to Benzoyl Peroxide, Arch.Orthop.Traumat.Surg., 94, 175 ~ 178, 1979
(0)	
Туре	: Patch test
Results	: Positive
Remarks	: For investigation of the sensitizing activity, several series of patients were patch tested. 11 subjects were tested with 1 % benzoyl peroxide only; 9 were positive. 30 patients were tested with 1 %, 0.5 % and 0.25 %. In 16 cases, all levels produced positive reactions; only 8 patients in this group showed no positive reaction. 31 of 41 patients treated, who had been negative before treatment, showed positive reaction at least to benzoyl peroxide 1 %
Reliability	: (2) Reliable with restriction
References	: Agathos, M., Bandmann, H.J., Benzoyl peroxide contact allergy in leg ulcer patients, Contact Dermatitis, 17, 316 ~ 317,1984

(p) Type Results Remarks	 Patch test Positive Benzoyl peroxid who had particip who had used a or greater. 	e patcl bated i benzog	h testi n a do yl per	ng wa ouble- oxide	as per -blind prod	forme acne uct sh	ed on stud	60 y y. 25 l reac	oung % o tivity	adults f those v of +1	
		No. of subjects givi					ing r	ng response			
	Treatment	+2	+1	±	0	+2	+1	±	0	Tota 1	
	Vehicle	0	4	2	12	0	0	3	15	18	
	Salicylic acid	0	2	1	11	0	0	0	14	14	
	Benzoyl peroxide	1	3	5	6	1	0	0	14	15	
	Benzoyl-peroxide										
	Salicylic acid	2	1	1	9	0	0	0	13	13	
Reliability References	 clinical allergy a majority of individual containing benzieffects. (2) Reliable with Rietschel, R.L. an acne study group 	ind had viduals zoyl p n restri and D oup, C	d dran with eroxic ction uncan	ns in natica positi le da , S.H t Derr	lly po ve pa ily w ., Ber natitis	tch te vithou nzoyl s, 8, 3	patcl sts co t sig perox 23 ~ 1	h test buld u nifica xide 1 326,	s, wh se pr int a reacti 1982	ons in	
(q) Type Results Remarks Reliability References	 Patch test Positive (allergie A 60 years old sacristan suffered had had 'allergy veneer with a sp airborne allergen the upper neck Benzoyl peroxic All candles lit in and beeswax w peroxide and sha (2) Reliable with Bonnekoh, B. an from benzoyl per Dermatitis, 24, 3 	Patch test Positive (allergic contact dermatitis) A 60 years old former joiner having served for about 15 yr as a sacristan suffered from eczema on his face, neck and hands. He had had 'allergy of face' at the age of 22 yrs, when busy staining veneer with a special was preparation. There were no reactions to airborne allergens on prick testing. Patch testing was performed on the upper neck with candle waxes and burned candle wicks Benzoyl peroxide gave a strong crescendo reaction at 0.5 % pet All candles lit in cathedral were made from a 9:1 mix. of paraffir and beeswax with cotton wicks. Wax bleaching with benzoyl peroxide and shaping candles were performed for several hours. (2) Reliable with restriction Bonnekoh, B. and Merk, H.F., Airborne allergic contact dermatitis from benzoyl peroxide as a bleaching agent of candle wax, Contac Dermatitis. 24, 367 ~ 368, 1984						yr as a ds. He taining ions to ned on wicks. % pet. araffin enzoyl urs. matitis Contact			

6. REFERENCES

Agathos, M., Bandmann, H.J., Benzoyl peroxide contact allergy in leg ulcer patients, Contact Dermatitis, 17, 316~317, 1984

Akzo Nobel, The Netherlands, Effects of the Wate - Accomodated Fraction of Lucidol on the Growth of the Freshwater Green Alga *Pseudokirchneriella subcapitata*, 1989

Aldrich, Catalog handbook of fine chemicals, Milwaukee, WI, Aldrich chem. Co., 163, 2000 ~ 2001

Autoxidized Squalene and Linoleic Acid, and of Simpler peroxides, in Relation to Toxicity of Radiation, Biochem. J. $67, 551 \sim 558, 1957$

Baker, J., Material Safety Data Sheet (MSDS) Number, B1831, Effective Date, 11/02/01 by Mallinckrodt chemicals

Basketter, D.A, Gerberick, G.F., An interlaboratory evaluation of the Buehler test for the identification and classification of skin sensitizers, Contact Dermatitis, 35, 146 ~151, 1996

Biozzi, G., Skin Tumorigenesis by Initiators and Promoters of Different Chemical Structures in Lines of Mice Selectively Bred for Resistance (Car-R) or Susceptibility (Car-S) to Two-stage Skin Carcinogenesis, Int. J. Cancer 83 (3), 335 - 340, 1999

Bonnekoh, B. and Merk, H.F., Airborne allergic contact dermatitis from benzoyl peroxide as a bleaching agent of candle wax, Contact Dermatitis, 24, 367 ~ 368,1984

Bretherick, L., Ed. Hazards in the Chemical Laboratory, 3rd Ed, The Royal Society of Chemistry. London, 270 ~ 271, 1981

Bretherick, L., Handbook of Reactive Chemical Hazards, Butterworths, 4TH Ed, London, 868 ~ 869, 1990

Buckingham, J., Ed., Dictionary of Organic Compounds, 5th Ed and Supplements, Chapman and Hall. New York, Vol. 2, 1612 ~ 1613, 1982

Budavari, S. (ed.)., The Merck Index - An Encyclopedia of Chemicals, Drugs, and Biologicals, Whitehouse Station, NJ: Merck and Co., Inc., on CD-ROM version 12:2, 10th edition, 1998

Cartwright, R.A., Hughes, B.R. and Cunliffe, W.J., Malignant Melanoma, Benzoyl Peroxide and Acne: A Pilot Epidemiological Case-Control Investigation, British J, Derm., 118, 239 ~ 242, 1988

Chem. Inspect Test Inst, Biodegradation and Bioaccumulation Data of Existing Chemicals Based on the CSCL Japan, Published by Japan Chemical Industry Ecology-Toxicology & Information Center, ISBN 4-89074-101-1,1992

Consumer Healthcare Products Association (CHPA), Dermal Oncogenicity Study of Benzoyl Peroxide Gels in Mice, Covance study No.6711-100, Vol 1 of 7, 2001

Dillon, D., Combes, R. and Zeiger, E., The Effectiveness of Salmonella strains TA100, TA102 and

TA104 for Detecting Mutagenicity of Some Aldehydes and Peroxide, Mutagenesis, 13 (1), $19 \sim 29$, 1998

Eaglstein, W.H., Allergic Contact Dermatitis to Benzoyl Peroxide, Arch. Derm., 97, p527, 1968

Eastman Kodak Co, laboratory of Industrial Medicine, Health, Safety, and Human Factors laboratory, 1964, 2 pp cited in 'criteria for recommended standard Occupational Exposure to Benzoyl Peroxide, NIOSH 1977 published'

Epstein, J., Photocarcinogenesis Promotion Studies with Benzoyl Peroxide and Croton Oil, J. Investigative Dermatology, 91(2), $114 \sim 116$, 1988

Epstein, S.S., Arnold, E., Andrea, J., Bass, W. and Bishop, Y., Detection of Chemical Mutagens by the Dominant Lethal Assay in the Mouse, Toxicology and Applied Pharmacology, 23, 288 ~ 325, 1972

Fujita, H. and Sasaki, M., Mutagenicity Test of Food Additives with *Salmonella typhimurium* TA 97 and TA102 (II), Annu. Rep. Tokyo Metro. Res. Lab.P.H., 38, 423 ~ 430, 1987 (in Japanese with English abstract)

Gebhardt, M., Geier, J., Evaluation of patch test results with denture material series, Contact Dermatitis, 34, 191 ~ 195, 1996

Gigiena Sanitariya. V/O Mezhdunarodnaya Kniga 113095 Moscow, USSR Vol.1-, 1936 cited in SAX'S Dangerous Properties of Industrial Materials, Tenth Edition, 1999 John Wiley & Sons, Inc CD version

Gimenez-Conti, I.B., Oubari, D., and Slaga, T.J., Dose-response of benzoyl peroxide hyperplastic promoter and possible complete carcinogenic activities, Proc. Am. Assoc., Cancer Res., 34, 184, 1993

Ginkel, C.G. van, Akzo Chemicals BV, The Netherlands, Biodegradability of Dibenzoyl peroxide, 1990

Ginkel, C.G. van and Stroo, C.A., Akzo Chemicals BV, The Netherlands, Toxicity of dibenzoyl peroxide for activated sludge, 1990

Haustein, U.F., Kramer, I., Fickweiler, E., Beahandlung der Akne vulgaris mittels Benzoylperoxid, Dermatol. Monatsschr., 167, 149 ~153, 1981

Haustein, U-F, Tegetmeyer, L. and Ziegler, V., Allergic and Irritant Potential of Benzoyl Peroxide, Contact Dermatitis, 13, 252 ~ 257, 1985

Hazardous Chemical Database, hardy Research Group the University of AKRON, Department of Chemistry, 2001

Hogan, D.J., To T, Wilson, E.R., Miller, A.B., Robson, D., Holfeld, K., Lane, P., A study of acne treatments as risk factors for skin cancer of the head and neck. Brit. J. Dermatol., 125, 343 ~ 348, 1991

Hogan, D., Wilson, E., Miller, A., Benzoyl peroxide and facial skin cancer, Am. J. Epidemiol., 134, 772, Criteria for Recommended standard, Occupational Exposure to Benzoyl Peroxide, NIOSH, 1977

OECD SIDS	BENZOYL PEROXIDE
6. REFERENCES	ID 94-36-0
	DATE: AUGUST 2002

Horgan, V.J., Philpot, J. ST L., Porter, B.W. and Roodyn, D.B., Toxicity of Autoxidized Squalene and Linoleic Acid, and of Simpler peroxides, in Relation to Toxicity of Radiation, Biochem. J., 67, 551-558, 1957

International Agency for Research on Cancer, World Health Organization, IARC Monographs on the Evaluation of Carcinogenic Risk of Chemicals to Man, International Agency for Research on Cancer. Geneva, Vol. 36, 267 ~ 283, Supplement 7, p. 58.

Ishidate, M.J., Yoshikawa, K. and Soluni, T., Mutagenicity Testing of Food Additives-First Screening Data Hegen-to-dokusei, Mutagens Toxicol., 3, 82 ~ 90, 1980

Iversen, O.H., Skin Tumorigenesis and Carcinogenesis Studies with 7,12dimethylbenzene[a]anthracene, ultraviolet Light, Benzoyl Peroxide (Panoxy gel 5 %) and Ointment Gel Carcinogenesis, 9 (5), $803 \sim 809$, 1988

Jager, M. and Balda, B.R., Loosening of a Total Hip Prosthesis at Contact Allergy Due to Benzoyl Peroxide, Arch. Orthop.Traumat. Surg., 94, 175 ~ 178, 1979

Kimber, I., Hilton, J., Dearman, R. J., Gerberick, G. F., Ryan, C. A., Basketter, D. A., Lea, L., House, R. V., Ladics, G. S. Loveless, S. E. and Hastings, K. L., Assessment of the Skin Sensitization Potential of Topical Medicaments Using the Local Lymph Node Assay: An Interlaboratory Evaluation, J. Toxicol. Environ. Hlth., Part A, 53 (7), 563 ~ 579, 1998

Kligman, A.M. and Kligman, L.H., A Hairless Mouse Model for Assessing the Chronic Toxicity of Topically Applied Chemicals, Food and Chemical Toxicology 36, 867 ~ 878, 1998

Korhonen, A., Hemminki, K. and Vainio, H., Embryotoxic Effects of Eight Organic Peroxides and Hydrogen Peroxide on Three-Day Chicken Embryos, *Environ. Res.*33, 54 ~ 61, 1984

Kuchle, H.J., Untersuchungen uber die augenschadigende Wirkung organischer Peroxyde, Zentralblatt fur Arbeitsmedizin und Arbeitsschuntz, Heit, 2, 25 ~ 31, 1958

Kurokawa, Y., Takamura, N., Matsushima, Y., Imazawa, T. and Hayashi, Y., Studies on the Promoting and Complete Carcinogenic Activities of Some Oxidizing Chemicals in Skin Carcinogenesis, Cancer Letters, 24, 299 ~ 304, 1984

Lewis, R.J., Sr., Eds, Hawley's Condensed Chemical Dictionary, 14th Ed John Wiley & Sons, Inc. New York, 127 ~ 128, 2001

Leyden, J.J., Kligman, A.M, Contact sensitization to benzoyl peroxide, Contact Dermatitis, Vol. 3, 273 ~ 275, 1977

Lide, D.R. (ed.), CRC Handbook of Chemistry and Physics, 76th ed, Boca Raton, FL, CRC Press Inc., 3 ~ 250, 1995 ~1996

Lyman, W.J. *et al*, Handbook of Chemical Property Estimation Methods, Washington DC, Amer. Chem. Soc., $15 \sim 1$ to $15 \sim 29$, 1990

Marhold, J.V., Sbornik Vysledku Toxixologicheho Vysetreni Latek A Pripravku, Institut Pro Vychovu Vedoucicn Pracovniku Chemickeho Prumyclu Praha, Czechoslovakia, 1972. cited in SAX'S Dangerous Properties of Industrial Materials, Tenth Edition, 1999 John Wiley & Sons, Inc CD version

Mark, U., Meuwsen, I.J.B., Akzo Chemicals BV, The Netherlands, Acute Toxicity of Dibenzoyl peroxide to fish, 1989

Meylan, W.M. and Howard, P.H., Bond contribution method for estimating Henry's law constants, Environ. Toxicol. Chem, 10, 1283 ~ 1293, 1991

Morillas, I.M., Martinesz, A.ALozano, J.L.S. and Perez, A.G., Is benzoyl peroxide an irritant or sensitizer?, Contact Dermatitis, 16, 232 ~ 233,1987

Morsches, B. and Holzmann H., Untersuchungen zur perkutanen Resorption von Benzoylperoxid, Arzneim. -Forsch. / Drug Res., 32 (1), Nr.3, 1982

Mosby Year Book, Contact and Occupational Dermatology, St. Louis, MO/Marks, J.G., V.A. DeLeo, 164 ~ 164, 1992

Nacht, S., Yeung, D., Joseph, N., Beasley, J. N. Jr., Anjo, M. D. and Maibach, H.I., Benzoyl Peroxide: Percutaneous Penetration and Metabolic Disposition, J. Am. Acad. Dermatol., 4 (1), $31 \sim 37$, 1981

National Fire Protection Association, Fire Protection Guide on Hazardous Materials, 7th Ed, National Fire Protection Association.Quincy, MA, 49~113, 49~114, 491M~57, 491M~145, 1981

National Institute of Environmental Research (NIER), Korea, survey on circulation volume and use pattern of benzoyl peroxide in Korea, 2001

National Institute of Environmental Research (NIER), Korea, Estimation of Physical Chemical Properties and Environmental Fate of SIDS Chemicals (II)-on Benzoyl peroxide, 2002

National Institute of Environmental Research (NIER), Korea, "Hydrolysis as a Function of pH" (tested by LGCI), 2001

National Institute of Environmental Research (NIER), Acute Eye Irritation/Corrosion Test in Rabbits (Report No. S367 tested by LGCI), 2002

National Institute of Environmental Research (NIER), Acute oral toxicity of benzoyl peroxide (Report No. G01083 tested by KRICT), 2001

National Institute of Environmental Research (NIER), Combined Repeated Dose Toxicity with the Reproduction /Developmental Toxicity Screening Testing of Benzoyl peroxide in Rats (Report No.P049 tested by LGCI), 2001

National Institute of Environmental Research (NIER), Korea, Bacterial Reverse Mutation Test (Test No. S350 tested by LGCI), 2001

National Institute of Environmental Research (NIER), Korea; Micronucleus test of Benzoyl peroxide in mouse (Test No.S349 tested by LGCI), 2001

National Institute of Environmental Research (NIER), Korea, The acute toxicity of Benzoyl peroxide

to Aquatic Invertebrate (Daphnia) (Report No. EG01006 tested by KRICT), 2001

National Institute of Environmental Research (NIER), Korea, The toxicity of Benzoyl peroxide to Aquatic plants (algae) (Report No. EG01005, tested by KRICT), 2001

National Institute of Environmental Research, Korea (NIER), The toxicity of Benzoyl peroxide to earthworm (tested by KRICT), 2002

National Institute of Environmental Research (NIER), Korea, The acute toxicity of Benzoyl peroxide to fish (Report No. EG01077 tested by KRICT), 2002

NIOSH, Criteria for Recommended Standard, Occupational Exposure to Benzoyl Peroxide, 1977

O'Connell, J.F., Klein-Szanto, A.J.P., DiGiovanni, D.M., Fries, J.W. and Slaga, T.J., Enhanced Malignant Progression of Mouse Skin Tumors by the Free-Radical Generator Benzoyl Peroxide, Cancer Research, 46 (6), 2863 ~ 2865, 1986

Odukoya, O. and Shklar, G., Initiation and Promotion in Experimental Oral Carcinogenesis, Oral Surg. Oral Med. Oral Pathol., 58 (3), 315 ~ 320, 1984

OECD, Environment Directorate, Joint Meeting of the Chemicals Committee and the Working Party on Chemicals, Pesticides and Biotechnology, OECD Series on Testing and Assessment Number 27, Guidance Document on the Use of the Harmonised System for the Classification of Chemicals Which are Hazardous for the Aquatic Environment, ENV/JM./MONO(2001)8

Online National Toxicology Program (NTP), 2002

Online Toxicology Data Network (TOXNET), Hazardous Substances Data Bank (HSDB), 2002

Philpot, J. St. L. and Roodyn, D.B., A Comparison Between the Effects in Mice of Injected Organic Peroxides and of Whole-body X-Irradiation, Int. J. Rad. Biol. 4, 372 ~ 382, 1959

Richard, J., Lewis, Sr., SAX' Dangerous Properties of Industrial Materials, John Wiley & Sons, Inc. New York, on CD-ROM, 10th Ed, 2000

Rietschel, R.L. and Duncan, S.H., Benzoyl peroxide reactions in an acne study group, Contact Dermatitis, 8, $323 \sim 326$, 1982

Rezazadeh, H. and Athar, M., Effect of Iron Overload on the Benzoyl Peroxide Mediated Tumor Promotion in Mouse Skin, Cancer Letters 126, 135 ~ 142, 1998

Saladino, A.J., Willey, J.C., Lechner, J.F., Grafstrom, R.C., LaVeck, M. and Harris, C.C., Effects of Formaldehyde, Acetaldehyde, Benzoyl Peroxide, and Hydrogen Peroxide on Cultured Normal Human Bronchial Epithelial Cells, Cancer Research, 45, 2522 ~ 2526, 1985

Sangster J, LOGKOW Databank, Sangster Res. Lab., Montreal Quebec, Canada, 1994

Santiago Q., Olaguibel, J.M., Garcia, B.E. and Tabar, A.I., Occupational airborne contact dermatitis due to benzoyl peroxide, Contact Dermatitis. 29, 165 ~ 166, 1993

Saran, A., Pazzaglia, S., Rebessi, S., Bouthillier, Y., Pioli, C., Covelli, V., Moutoin, D., Doria, G.
and Schweizer, J., Loehrke, H., Edler, L. and Goerttler, K., Benzoyl Peroxide promotes the Formation of Melanotic Tumors in the Skin of 7, 12-dimethylbenz[a]anthracene-initiated Syrian Golden Hamsters, Carcinogenesis, 8 (3), 479 ~ 482, 1987

Sevila A., Romaguera, C., Vilaplana, J., Botella, R., Contact dermatitis in children, Contact Dermatitis, 30, 292 ~ 294, 1994

Sharratt, M., Frazer, A.C. and Forbes, O.C., Study of the Biological Effects of benzoyl Peroxide, Fd Cosmet. Toicol. Vol.2, pp 527 ~ 538, 1964

Slaga, T.J., Klein-Szanto, A.J.P., Triplett, L.L.I., Yotti, L.P. and Trosko, J.E., Skin Tumor-Promoting Activity of Benzoyl Peroxide, a Widely Used Free Radical-Generating Compound, Science 213, 1023 ~ 1025, 1981

Swann, R.L. et al, Res Rev 85 23, 1983

Takahashi, A., Problems of Hygiene Maintenance for Food Coming into Contact with Rubber and Plastic Products, Nippon Gomu Kyokaishi 48 (9), 537, 1975 [Translated by English, E. A., Internat'l Polymer Sci. Technol., 3 (1), 93 ~ 105, 1976

The OECD HPV program, SIDS Initial Assessment Report for 13th SIAM- Benzoates category

Tkach, J.R., Allergic Contact Urticaria to Benzoyl Peroxide, Cutis 29, 187 ~ 188, 1982

To T, Hogan, D, Wilson, E, Miller, A, Benzoyl peroxide and facial skin cancer, Am. J. Epidemiol., 134, 772

Toxicity and health hazard summary, report No. 900713. Rochester, NY, Eastman Kodak Co, lab of ind. Med. Health, Safety&Human Factors Lab., 1964, 2p. cited in 'criteria for recommended standard Occupational Exposure to Benzoyl Peroxide, NIOSH 1977 published'

Von Bahmer, F.A., Schulze-Dirks, A. und Zaun, H., Sensibilisierende Wirkung einer fur die behandlung des Ulcus cruris verwendeten 20 % igen Benzoylperoxid-Zubereitung, Dermatosen, 32, Nr 1, $21 \sim 24$, 1984

U.S. Department of Health and Human Services and U.S. Department of Labor, NIOSH/OSHA Occupational Health Guidelines for Chemical Hazards, Vol. 3, DHHS (NIOSH) Publication No. 81 \sim 123, January, Volume 1, 1981

Wazeter, F.X. and Goldenthal, E., Acute Oral Toxicity Test in Male Albino Rats; International Research and Development Corporation (IRDC), August 31, 1973

Wazeter, F.X. and Goldenthal, E., Acute Inhalation to Rats; 328-005 International Research and Development Corporation (IRDC) August 31, 1973

Wazeter, F.X. and Goldenthal, E., Eye irritation in Albino Rabbits, International Research and Development Corporation (IRDC) August 31, 1973

Wazeter, F.X., and Goldenthal, E., Primary Skin Irritation and Corrosive Hazard Test in Albino Rabbits, International Research and Development Corporation (IRDC) August 31, 1973

Weinberg, J.M., Moss, T., Gupta, S.M., White, S.M. and Don, P.C., Reticulate Hyperpigmentation

of the Skin After Topical Application of Benzoyl Peroxide, Acta. Derm. Venereol (Stockh)., 78, 301 ~ 302, 1998

Wepierre, J., Corroller, M. and Didry, J. R., Distribution and Dissociation of Benzoyl Peroxide in Cutaneous Tissue after Application on Skin in the Hairless Rat, International Journal of Cosmetic Science $8, 97 \sim 104, 1986$

Wester, R.C., Patel, R., Nacht, S., Leyden, j., Melendres, J. and Maibach, H., Controlled Release of Benzoyl Peroxide from a Porous Microsphere Polymeric System Can Reduce Topical Irritancy, J. Am. Acad. Dermatol., 24, 720 ~ 726, 1991

Wijk, R.J., van and Groenenboom Ir.C.J., Akzo Nobel, The Netherlands, Acute Toxicity of Lucidol to *Daphnia magna*, 1999

Yeung, D., Nacht, S., Bucks, D. and Maibach, H. I., Benzoyl Peroxide: Percutaneous Penetration and Metabolic Disposition. II. Effect of Concentration, J. Am. Acad. Dermatol. 9, 920 ~ 924, 1983

Zeiger, E., Anderson, B., Haworth, S., Lawlor, T., Mortelmans, K. *Salmonella* Mutagenicity Tests: IV. Results from the Testing of 300 Chemicals. Env. Mol. Mutag. 11/12, 1 ~ 158, 1988