SIDS INITIAL ASSESSMENT PROFILE

CAS No.	56-81-5
Chemical Name	1,2,3-Propanetriol (Glycerol)
Structural Formula	НО ОН ОН
RECOMMENDATIONS	

The chemical is currently of low priority for further work.

SUMMARY CONCLUSIONS OF THE SIAR

Human Health

All SIDS health endpoints are fulfilled. It should be noted that much of the data on glycerol is historic and of rather low quality compared to current guideline requirements. Nevertheless, there is an overall consistency within the available data that allows conclusions to be drawn. Glycerol is absorbed following ingestion and metabolised by glycerokinase in the liver to carbon dioxide and water or incorporated in the standard metabolic pathways to form glucose and glycogen. The weight of evidence indicates that glycerol is of low toxicity when ingested, inhaled or in contact with the skin.

Glycerol is of a low order of acute oral and dermal toxicity with LD_{50} values in excess of 4000 mg/kg bw. At very high dose levels, the signs of toxicity include tremor and hyperaemia of the gastro-intestinal -tract. Skin and eye irritation studies indicate that glycerol has low potential to irritate the skin and the eye. The available human and animal data, together with the very widespread potential for exposure and the absence of case reports of sensitisation, indicate that glycerol is not a skin sensitiser.

Repeated oral exposure to glycerol does not induce adverse effects other than local irritation of the gastro-intestinal tract. The 2-year study of Hine (1953) was chosen to establish the overall NOEL after prolonged treatment with glycerol of 10,000 mg/kg bw/day (20% in diet), which is in agreement with the findings in other studies. At this dose level no systemic or local effects were observed. For inhalation exposure to aerosols, the NOAEC for local irritation effects to the upper respiratory tract is 165 mg/m3 and 662 mg/m3 for systemic effects.

Glycerol is free from structural alerts, which raise concern for mutagenicity. Glycerol does not induce gene mutations in bacterial strains, chromosomal effects in mammalian cells or primary DNA damage *in vitro*. Results of a limited gene mutation test in mammalian cells were of uncertain biological relevance. *In vivo*, glycerol produced no statistically significant effect in a chromosome aberrations and dominant lethal study. However, the limited details provided and the absence of a positive control, prevent any reliable conclusions to be drawn from the *in vivo* data. Overall, glycerol is not considered to possess genotoxic potential.

The experimental data from a limited 2 year dietary study in the rat does not provide any basis for concerns in relation to carcinogenicity. Data from non-guideline studies designed to investigate tumour promotion activity in male mice suggest that oral administration of glycerol up to 20 weeks had a weak promotion effect on the incidence of tumour formation.

No effects on fertility and reproductive performance were observed in a two generation study with glycerol administered by gavage (NOAEL 2000 mg/kg bw/day). No maternal toxicity or teratogenic effects were seen in the rat, mouse or rabbit at the highest dose levels tested in a guideline comparable teratogenicity study (NOEL 1180 mg/kg bw/day).

Environment

All SIDS environmental endpoints are fulfilled. It should be noted that much of the data on glycerol is historic and of rather low quality compared to current guideline requirements. However, the weight of evidence indicates that glycerol is of low toxicity to aquatic organisms and this conclusion is supported by QSAR predictions. The lowest LC50 for fish is a 24-h LC₅₀ of >5000 mg/l for *Carassius auratus* (Goldfish) and for aquatic invertebrates, a 24h EC50 of >10000 mg/l for *Daphnia magna* is the lowest EC50. Several tests on algae are available, which suggest very low toxicity to a range of species, however their validity is uncertain. A QSAR prediction for the 96h EC50 to algae was 78000 mg/l. No toxicity towards the microorganism *Pseudomonas putida* was observed at 10000 mg/l after exposure for 16 hours. No long-term aquatic toxicity data is available. Screening studies are available on frog and carp embryos which indicate some effects on growth and hatching rates respectively at very high concentrations of glycerol, >7000 mg/l. However, their ecological relevance is not clear.

In view of the limited robustness of the studies present, it was decided to derive a tentative PNEC for aquatic organisms using QSAR predictions of acute toxicity. The tentative PNEC for aquatic organisms is calculated to be 780 mg/L, based on the lowest QSAR value (calculated for algae EC_{50} 77,712 mg/L) and applying an assessment factor of 100 in accordance with the OECD guidance. An assessment factor of 1000 for the aquatic PNEC compartment could also be considered to reflect the uncertainty in the use of QSAR-predicted values. There are no sediment or terrestrial effect data, but partitioning to both soil and sediment is expected to be very low, based on the very low log K_{ow} of glycerol. The equilibrium partitioning method was used to calculated tentative PNECs for soil and sediment based on the PNEC_{aquatic} of 777 mg/l, PNEC_{sediment} = 479 mg/kg wwt and PNEC_{soil} = 92.1 mg/kg wwt.

Exposure

The worldwide market for glycerol for the year 2000 was 500,000 tonnes. Glycerol has widespread use and can be found in industrial, professional and consumer products. Glycerol is used as a constituent in numerous products and as an intermediate in industrial applications for the manufacture of products such as soaps/detergents and glycerol esters. It is found in consumer products such as pharmaceuticals, cosmetics, tobacco, food and drinks and is present in numerous other products such as paints, resins and paper.

There is a potential for occupational exposure through inhalation and skin contact. Consumers may be exposed to glycerol by the oral and dermal routes of exposure. Smoking may lead to an additional glycerol uptake by inhalation.

There is potential exposure to the aquatic compartment arising from the production and processing of this substance. Glycerol will enter the aqueous and terrestrial environment from end uses such as in consumer products and down hole lubricants for oil and gas fields.

Glycerol is a liquid with a calculated vapour pressure of 0.000106 hPa (at 25°C), is fully miscible with water and has a Log K_{ow} of -1.76 (measured). It has a calculated half-life for photo-oxidation of \sim 7 hours and is not susceptible to hydrolysis. The experimental data indicate that glycerol is readily biodegradable under aerobic conditions. Fugacity modelling (Mackay Level III) predicts that glycerol will partition to the aquatic compartment (100%). Based on the low Log Kow, it has a low potential for sorption to soil and is not expected to bioaccumulate.

NATURE OF FURTHER WORK RECOMMENDED

No further work is indicated, because of the low hazard potential of this substance.